



1. Home (<https://www.gov.uk/>)
  2. Coronavirus (COVID-19) (<https://www.gov.uk/coronavirus-taxon>)
  3. Healthcare workers, carers and care settings during coronavirus (<https://www.gov.uk/coronavirus-taxon/healthcare-workers-carers-and-care-settings>)
  4. Supported living services during coronavirus (COVID-19) (<https://www.gov.uk/government/publications/supported-living-services-during-coronavirus-covid-19>)
- Department of Health & Social Care (<https://www.gov.uk/government/organisations/department-of-health-and-social-care>)
  - Public Health England (<https://www.gov.uk/government/organisations/public-health-england>)

Guidance

# COVID-19: guidance for supported living

Updated 17 May 2021

## Contents

Who this guidance is for

Steps that supported living providers and local authorities can take to maintain service delivery

Risk assessment, risk reduction and local implementation

Staff within clinically vulnerable groups

General infection prevention and control

Visits in and out of supported living settings

If a supported living worker has COVID-19 symptoms

If someone in supported living has symptoms of COVID-19

Annex A: taking swabs

Annex B: additional resources

[Print this page](#)



© Crown copyright 2021

This publication is licensed under the terms of the Open Government Licence v3.0 except where otherwise stated. To view this licence, visit [nationalarchives.gov.uk/doc/open-government-licence/version/3](https://www.nationalarchives.gov.uk/doc/open-government-licence/version/3) (<https://www.nationalarchives.gov.uk/doc/open-government-licence/version/3>) or write to the Information Policy Team, The National Archives, Kew, London TW9 4DU, or email: [psi@nationalarchives.gov.uk](mailto:psi@nationalarchives.gov.uk).

Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.

This publication is available at <https://www.gov.uk/government/publications/supported-living-services-during-coronavirus-covid-19/covid-19-guidance-for-supported-living>

## Who this guidance is for

This guidance is designed to update and build on the previous advice to supported living providers, which was withdrawn on 13 May 2020. It sets out:

- key messages to assist with planning and preparation in the context of the coronavirus (COVID-19) pandemic so that local procedures can be put in place to minimise risk and provide the best possible support to people in supported living settings. These local procedures may need to be updated to reflect changes in government guidance and advice as the pandemic response changes
- safe systems of working, including social distancing, respiratory and hand hygiene and enhanced cleaning
- how infection prevention and control (IPC) and personal protective equipment (PPE) (<https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control>) applies to supported living settings

Although it is primarily intended for the managers, care and support workers, and other staff in supported living settings, it is also relevant to local authorities, clinical commissioning groups (CCGs), primary care networks (PCNs), and community health services. It will also be useful to read this guidance alongside the materials listed in Annex B.

Supported living enables adults to live in their own home – with the help they need to be independent – and allows them to choose where they want to live, who with, how they want to be supported, and what happens in their own home. As a term, it covers a wide variety of settings and may include some form of group living.

Homes may be shared between several people and have communal space or consist of separate units of self-contained accommodation – with or without communal space – but which may be located in shared buildings such as a block of flats and/or on shared grounds. Elements of supported living may be formally regulated by Care Quality Commission (CQC) and others not.

Supported living services involve tenure rights – renting or ownership, with associated occupancy rights. Some provide regulated ‘personal care’ and others support daily living activities such as help with shopping, food preparation, access to the community or a combination of both.

In some supported living models, it is not possible to defer the care and/or support provided to another day without putting people at risk of harm. It is therefore vital that these services are maintained.

The guidance is primarily for supported living settings, but many of the principles are applicable to extra care housing for older people. It may also be a useful resource for the wider supported housing sector, such as retirement or sheltered housing. Given the different types of supported living and the associated care, support and help for people living there, this guidance cannot be specific to individual locations, and local managers should use it to develop their own specific ways of working to protect people’s wellbeing and minimise risks.

Although this guidance is often worded as if the organisation’s management has full responsibility for accommodation as well as support and care services, ordinarily, care providers have no responsibility for property, accommodation or environment issues in supported living. In many instances, management’s

role will be to develop local procedures and work with the people being supported and, with consent, their families, GP, support groups, and care/support providers to ensure that individual plans are in place to protect wellbeing and minimise risks.

In some instances, home modifications or adaptations may be required, for example to receive deliveries or to support reablement. Supported living is the person's own home and, although advice and good practice may be offered, it will only be followed if the person understands the advice and is in agreement.

Some people being supported may lack capacity to understand and make decisions based on advice about the COVID-19 pandemic. It is important that all steps are taken to communicate information with people in a way that they are most likely to be able to understand. For example, autistic people and people with learning disabilities, dementia, or mental ill health may have difficulties with understanding complex instructions or forget them. This, and the other principles and requirements of the Mental Capacity Act 2005 (<https://www.scie.org.uk/mca/introduction/mental-capacity-act-2005-at-a-glance>) (MCA) must be followed when it is felt a person being supported may lack capacity. Guidance (<https://www.gov.uk/government/publications/coronavirus-covid-19-looking-after-people-who-lack-mental-capacity/the-mental-capacity-act-2005-mca-and-deprivation-of-liberty-safeguards-dols-during-the-coronavirus-covid-19-pandemic#use-of-the-mca-and-dols-due-to-covid-19>) and additional advice is available about the application of the MCA during the pandemic (<https://www.gov.uk/government/publications/coronavirus-covid-19-looking-after-people-who-lack-mental-capacity/the-mental-capacity-act-2005-mca-and-deprivation-of-liberty-safeguards-dols-during-the-coronavirus-covid-19-pandemic-additional-guidance#other-settings>).

## **Steps that supported living providers and local authorities can take to maintain service delivery**

To maintain service delivery, supported living providers and local authorities are advised to follow these steps.

### **1. Ensure that lists of people in supported living are up to date**

Supported living providers and local authorities should work together to ensure that, where feasible, their lists of people in supported living are up to date. Lists should establish the levels of formal and informal care and support available to individuals.

Individuals should be supported to draw up a contingency plan – with their care providers and any unpaid carers – that can be enacted should they contract COVID-19 or there be an impact on care delivery due to COVID-19. Providers should consider how they could share this information.

### **2. Business continuity planning**

Providers should have business continuity plans to help them to manage in emergency situations. These should be kept up-to-date and key details to record may include:

- who provides care for the people in supported living environments and whether these individuals are still able to provide care and are not shielding/self-isolating, whether paid staff or informal carers
- how and where care and support plans are located
- requirements for any specialist care and/or long-term conditions

- key contacts coordinating care from other community-based services including, but not limited to, mental health, learning disabilities, dementia, third sector Voluntary Social and Community Enterprises (VSCs), drug and alcohol or social work teams, and family members

### **3. Key contingency details**

Key contingency details should also include information about the person's modes of communication including technology, their likely reaction to changes in routine or unfamiliar carers, and ways to reduce potential stress. In cases where current circumstances make consistency impossible, providers should prepare people for the fact that it may be necessary for a different carer to support them.

It is particularly important to ensure risk management plans are updated for people who may find any change in routine challenging, for example autistic people or people living with dementia.

### **4. Mutual aid, care and support plans**

Providers and local authorities should work together to facilitate mutual aid, care and support plans across their areas. This is to inform planning ahead of a possible outbreak. Useful resources can be found on the Local Government Association website (<https://www.local.gov.uk/our-support/coronavirus-information-councils/covid-19-service-information/covid-19-public-health>).

### **5. Identify people who use direct payments or fund their own support**

Providers and local authorities should also work together to identify people who use direct payments or who fund their own support and help them establish the levels of support available from other providers or individuals.

It may be helpful for providers to share the number of hours of care they provide to help with planning, but they will want to satisfy themselves that it is lawful for them to share that information and get consent from the person where possible.

### **6. Avoid sharing staff between settings**

Sharing staff between settings should be avoided to reduce the potential spread of COVID-19 from one setting to another. If a local risk assessment identifies service delivery issues caused by low staffing, then supported living and care/support providers can work with local authorities to establish plans for mutual aid, including limited sharing of the workforce.

Local primary and community health services providers may support with the deployment of volunteers and agency staff where that is safe to do so and provided safeguarding measures are in place.

### **7. Identify people who are clinically extremely vulnerable**

Identify people who are clinically extremely vulnerable (<https://www.gov.uk/government/publications/guidance-on-shielding-and-protecting-extremely-vulnerable-persons-from-covid-19>), and work with them, their families or advocates to explain issues related to guidance and make a joint decision on how they will be supported and on their accommodation and support needs. For example, when shielding advice is operational, a person who uses services may want to remain in their current home if they can be supported to 'shield' or they may wish/need to move to different accommodation that will enable them to 'shield' effectively.

## 8. Maintain oversight of people who are self-isolating

The supported living provider should maintain oversight of people who are self-isolating, and note the arrangements that local authorities, CCGs and NHS 111 are putting in place to refer people self-isolating at home to volunteers who can offer practical and emotional support.

By following these steps, most people who live in supported living environments should have a continuity of care, support and help that adapts to the situation with COVID-19. For a small number of people, where a person's wellbeing is at risk, the managers of supported living environments may wish to contact social workers in their local authority to seek further advice and support.

### Risk assessment, risk reduction and local implementation

A suite of guidance (<https://www.gov.uk/coronavirus>) including responses to frequently asked questions (<https://www.gov.uk/government/publications/coronavirus-outbreak-faqs-what-you-can-and-cant-do/coronavirus-outbreak-faqs-what-you-can-and-cant-do>) has been published to support people in making decisions related to coronavirus (COVID-19) and many of these resources will be relevant to the supported living sector.

Providers will need to consider how to implement and risk assess these recommendations according to their individual circumstances and local operating models and may also wish to refer to JCC and PPE recommendations for:

- how to work safely in care homes (<https://www.gov.uk/government/publications/covid-19-how-to-work-safely-in-care-homes>)
- how to work safely in domiciliary care (<https://www.gov.uk/government/publications/covid-19-how-to-work-safely-in-domiciliary-care>)
- admission and care of people in care homes (<https://www.gov.uk/government/publications/coronavirus-covid-19-admission-and-care-of-people-in-care-homes>)

Updated guidance for homeless hostels has been developed (<https://www.gov.uk/government/publications/covid-19-guidance-on-services-for-people-experiencing-rough-sleeping>). There is also guidance issued by Public Health England (PHE) for individuals, families and informal care workers for households with possible coronavirus infection (<https://www.gov.uk/government/publications/covid-19-stay-at-home-guidance>).

Please note that this guidance is of a general nature and an employer should consider the specific conditions of each individual place of work and comply with all applicable legislation.

### Staff within clinically vulnerable groups

Supported living settings are staffed by a wide range of people and some may be more vulnerable to infection, for example, because they have an underlying health condition. Staff whose health makes them clinically extremely vulnerable are recommended to follow the guidance on shielding and protecting clinically extremely vulnerable persons from COVID-19 (<https://www.gov.uk/government/publications/guidance-on-shielding-and-protecting-extremely-vulnerable-persons-from-covid-19/guidance-on-shielding-and-protecting-extremely-vulnerable-persons-from-covid-19>).

Factors including age, sex, ethnicity, certain underlying health conditions (<https://www.gov.uk/government/publications/covid-19-review-of-disparities-in-risks-and-outcomes>) and/or pregnancy may be associated with an increased risk of or from COVID-19. Employers should ensure that

an appropriate person, such as a line manager, carries out individual conversations with all staff who may be at greater risk, in line with the latest guidance

(<https://www.gov.uk/government/publications/coronavirus-covid-19-reducing-risk-in-adult-social-care>).

Staff from black, Asian and minority ethnic (BAME) backgrounds may have increased concerns about COVID-19 (<https://www.gov.uk/government/publications/covid-19-understanding-the-impact-on-bame-communities>), and employers should handle these conversations sensitively. Employers should ensure that staff are supported and any necessary steps to reduce risk are considered on an individual and proportionate basis. The employee should consult their employer if they have any concerns and discuss issues raised with their line manager.

Many staff will be able to work normally, while being particularly careful to follow social distancing measures.

## General infection prevention and control

Infection prevention and control (IPC) measures include a hierarchy of controls designed to prevent harm and reduce transmission of infection to patients, residents, people who use supported living services, and health and social care staff and their co-workers.

To reduce the risk of COVID-19 spread and introduction into supported living facilities, measures should be followed including social distancing (<https://www.gov.uk/government/publications/staying-alert-and-safe-social-distancing/staying-alert-and-safe-social-distancing>), hygiene principles (hand hygiene, sneezing or coughing into a tissue, safe disposal of tissues, following environmental cleaning regimes (<https://www.gov.uk/government/publications/covid-19-decontamination-in-non-healthcare-settings>)) and self-isolation (if a person or member of their household becomes ill with COVID-19 symptoms).

In addition to these core measures, PPE (<https://www.gov.uk/government/publications/covid-19-how-to-work-safely-in-domiciliary-care>) is required in specific scenarios, when social distancing is not possible, as described in the PPE section of this guidance.

Staff providing care for autistic people and people with dementia or learning disabilities should make every effort to make sure that the people they support are aware of the key behaviours needed to follow good IPC and should provide encouragement and reminders when not followed. Staff should consider how the person they are supporting is most likely to understand the information and use the most appropriate communication techniques for that person.

This may include the use of social stories, information in pictorial form, engaging with friends and family members to support understanding and having regular online contact with the person being supported, when it is not possible to see them face to face, to reinforce IPC messaging. Learning Disability England (<https://www.learningdisabilityengland.org.uk/what-we-do/keeping-informed-and-in-touch-during-coronavirus/>) and PHE (<https://www.gov.uk/government/publications/covid-19-supporting-adults-with-learning-disabilities-and-autistic-adults/coronavirus-covid-19-guidance-for-care-staff-supporting-adults-with-learning-disabilities-and-autistic-adults#supporting-change>) have created resources which may support this messaging.

## Social distancing

The national advice on social distancing is to maintain 2 metres if at all possible. This is especially true in health and social care settings where the populations being cared for are vulnerable. Therefore, in this guidance document we recommend continuing to use 2 metres in all settings.

Whenever possible, staff should follow social distancing guidance

(<https://www.gov.uk/government/publications/staying-alert-and-safe-social-distancing>) being at least 2 metres away from the person they support. If this is not possible due to having to provide personal care, or if the person they support has behaviours and needs which make this difficult, then P.P.E.

([https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/886370/COVID-19\\_Infection\\_prevention\\_and\\_control\\_guidance\\_Appendix\\_2.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/886370/COVID-19_Infection_prevention_and_control_guidance_Appendix_2.pdf)) may be needed as highlighted below.

## Hand hygiene

Supported living staff should think carefully about ways that the person they support can be encouraged to participate in regular handwashing:

- washing hands with soap and water for at least 20 seconds is essential before and after all contact with the person, removal of protective clothing and cleaning of equipment and the environment
- where possible, promote hand hygiene and ensure that liquid soap and disposable paper towels are available at all sinks in shared areas
- alcohol-based hand rub can be used, where safe to do so, if hands are not visibly dirty or soiled and where appropriate, it should be accessible and have adequate provision
- if people you support are having visitors, encourage them to follow good respiratory and hand hygiene, washing their hands on arrival, during their stay, and on leaving (more details in visitors section)

## Respiratory and cough hygiene – ‘Catch it, bin it, kill it’

Disposable single-use tissues should be used to cover the nose and mouth when sneezing, coughing or wiping and blowing the nose. Used tissues should be disposed of promptly. Hands should be cleaned with soap and water, or alcohol hand rub where this is not possible and if safe, after coughing or sneezing, using tissues or after contact with respiratory secretions and/or contaminated objects.

Encourage individuals to keep hands away from the eyes, mouth and nose. Some people may need assistance with containment of respiratory secretions. Those who are immobile will need alcohol hand rub for hand hygiene, and a bag at hand for immediate disposal of the waste potentially contaminated with the COVID-19 virus, such as tissues. These bags should be placed into another bag, tied securely and kept separate from other waste. This should be put aside for at least 72 hours before being put in the usual household waste bin for disposal as normal. Where the person has learning disabilities, it will be important to make sure they understand exactly what they need to do and why.

## Visits in and out of supported living settings

Maintaining opportunities for visiting and spending time together is critical for the health and wellbeing of people being supported, and their relationships with friends and family. In addition, for many people, there are important reasons for having in-person visits, as not doing so may be difficult to understand and lead to distress.

There are risks that need to be considered – even where people are vaccinated – but these are risks that can be appropriately managed.



As stated above this guidance is intended for supported living settings, but many of the principles are applicable to extra care housing for older people. It may also be a useful resource for the wider supported housing sector, such as retirement or sheltered housing.

The approach described below for developing a policy and mitigating the risks of visits (both into and out of the home) has 3 key elements:

- people living in supported living settings live in their own homes and should be treated as such. This means they, and their visitors, need to follow the same national restrictions as other members of the public, including following each step in the government's roadmap around social contact. The roadmap and associated regulations provide some flexibilities which may apply to people in certain supported living settings (such as exemptions for some indoor gathering and in relation to forming support bubbles)
- supported living managers should seek to support and facilitate these opportunities wherever it is safe to do. They should develop policies for visits into and out of the setting, that are based on a dynamic risk assessment, and include consideration of the individual needs of the people who live there. These risk assessments should be developed in consultation with them
- supported living managers should also work with the people being supported to identify what further steps they can take in order to manage and mitigate risks that arise from visiting

The default position set out in this guidance is that visits should be supported and enabled wherever it is safe to do so.

## Following national restrictions

Some of the rules on what you can and cannot do (<https://www.gov.uk/guidance/covid-19-coronavirus-restrictions-what-you-can-and-cannot-do>) changed on 29 March. However, many restrictions remain in place. People in supported living and extra care settings, and those wishing to participate in visits with them, must follow any national restrictions in effect at the time of the visiting, just like any other member of the public.

The specific circumstances of the setting, and the people who live there, will also be relevant. Some people will be living as single person households, some as multiple person households where they share facilities such as kitchens, communal indoor areas and so on.

Together, these factors will largely define the range of visiting that is possible – both into and out of the setting. In most circumstances people living in supported living settings and extra care should be encouraged to take the opportunities presented by the lifting of restrictions to meet their friends and family – for example:

- from 8 March people can meet outdoors, for exercise or recreation outdoors (but only with their own household, support or childcare bubble, or with one other person from another household). This would enable someone to visit an outdoor space outside of the setting with one other person who is not already in their support bubble
- from 29 March people can meet outdoors (including in private gardens) with up to 6 people who are not in their household or support bubble, or more if limited to 2 households. This would enable someone to meet with others in a communal garden or outdoor space in the supported living or extra care setting, or equally an outdoor public space outside of the setting

- many people living in a supported living setting will be able to form a support bubble (<https://www.gov.uk/guidance/making-a-support-bubble-with-another-household>). For example, a single person household could form a bubble with another household (for example parents or family) and therefore receive visitors indoors at the setting, or for example make a visit to the family home

## Setting a visiting policy that supports these opportunities

Providers should seek to support visiting into and out of the setting as described in this guidance wherever it is possible to do so in a risk-managed way and in line with the principles set out below.

A wide range of professionals have an important role in supporting this and ensuring that visiting can happen safely, including scheme managers, Directors of Public Health (DPH) and Directors of Adult Services. Alongside these professionals, people being supported, relatives and families each have important roles to play.

In supported living environments the accommodation is the person's own home. However, it may also be a staff workplace and include a range of communal areas and shared facilities. It is therefore important to consider the risks arising from visits to those taking part, as well as the risks for others with whom they live, or may later come into contact including those supporting them, older relatives and so on.

Therefore, in all cases, arrangements for visiting into and out of the setting should be supported by a dynamic risk assessment for the overall setting, as well as an individualised assessment of the benefits of visiting and the risks to particular people because of their care and support needs. The risk assessment should consider people's rights as part of the wider roadmap and opening up of social activity. The risk assessment will also need to reflect whether the setting is a 'high risk' setting (as designated by the local DPH).

When planning a visit into or out of the setting, providers, and care and support workers should work with individuals and their families to consider their needs and maximise their safety. This will enable people being supported to make decisions about visits out of the home, and how these visits can be made possible.

## Advice for providers when taking visiting decisions for particular people

Any visiting arrangements should be made in agreement with the person being supported. If the person is assessed as not having capacity in relation to this decision, the provider should work within the appropriate MCA framework to establish whether the arrangements are in the person's best interests. The government has published advice on the MCA and application of Deprivation of Liberty Safeguards (DoLS) (<https://www.gov.uk/government/publications/coronavirus-covid-19-looking-after-people-who-lack-mental-capacity>) during the pandemic.

Regard should also be given to the ethical framework for adult social care (<https://www.gov.uk/government/publications/covid-19-ethical-framework-for-adult-social-care>), and the wellbeing duty in section 1 of the Care Act 2014 (<https://www.legislation.gov.uk/ukpga/2014/23/section/1/enacted>), and all decisions should be taken in light of general legal obligations, such as those under the Equality Act 2010 and Human Rights Act 1998, as applicable. Social workers can help providers to meet these duties by providing advice in individual cases should that be required.

Providers must consider the rights of people who may lack the relevant mental capacity needed to make a decision about visits out of their home. These people are protected by the empowering framework of the Mental Capacity Act (MCA) 2005 and its safeguards. The government has published advice on the

## **MCA and application of Deprivation of Liberty Safeguards (DoLS)**

(<https://www.gov.uk/government/publications/coronavirus-covid-19-looking-after-people-who-lack-mental-capacity/the-mental-capacity-act-2005-mca-and-deprivation-of-liberty-safeguards-dols-during-the-coronavirus-covid-19-pandemic-additional-guidance>) during the pandemic.

### **In the event of an outbreak**

If an outbreak is declared in the setting, the provider should take advice from the local health protection team who will undertake a risk assessment and determine subsequent next steps.

It might be the case that the local health protection team will recommend some visiting restrictions, these should continue until such time as it is understood that the outbreak has been brought under control.

### **Advice to help manage and mitigate risks for visits into the setting**

Managers should consider what further steps can be taken to mitigate risks related to the planned visit.

Visitor testing is a tool to help mitigate the risks of visiting. Testing is not a requirement for visiting and managers should not refuse visits to visitors who have not taken a test unless they are symptomatic or should be self-isolating for another reason (for example if they are a contact of someone who has tested positive). We recommend visitors participate in testing to reduce risk of introduction of infection through asymptotically infected people, in particular for higher risk settings with shared living accommodation spaces which have a higher potential for outbreaks.

Testing is one tool that can be used to help mitigate the risks to identify asymptomatic visitors who are likely to be infectious before they visit so they are able to self-isolate immediately. All providers who are currently eligible for staff testing (<https://www.gov.uk/government/publications/coronavirus-covid-19-testing-service-for-extra-care-and-supported-living-settings>) are able to access rapid lateral flow tests (LFTs) for the purpose of supporting safer visits. Managers can place an order for tests using their unique organisation number (UQN) from the test kit ordering portal. Each setting will receive 4 test kits per person per week which can be used to support both visits in and visits out (<https://www.gov.uk/government/publications/coronavirus-covid-19-lateral-flow-testing-in-adult-social-care-settings>).

Setting managers have discretion to set up their own testing areas with clinical guidance (<https://www.gov.uk/government/publications/coronavirus-covid-19-lateral-flow-testing-in-adult-social-care-settings>). Managers should ensure the testing area has enough space to allow visitors to maintain social distancing before, during and after the test, including a waiting area and a one-way system. The area should comply with fire safety regulations that govern deployment sites and hard, non-porous flooring that can withstand chlorine cleaning agents. Visitors should have ready access to hand hygiene and the area should be well ventilated with fresh air, either by appropriate ventilation systems or by opening windows and doors. Managers should also consider storage of tests.

Testing onsite at the setting is preferable for assurance purposes. However, recognising that individuals now have access to testing through other routes and visitors may be travelling long distances to visit, managers can allow visitors to provide evidence of a negative test taken on the day of the visit through other means including:

- assisted testing at another lateral flow site such as an asymptomatic testing site (ATS)
- self-testing at home through test kits provided by the setting once settings have access to packs of 7 test kits (which the **MHRA** has authorised for self-test use)

- self-testing at home using test kits provided by the government such as at a school, workplace, the universal testing offer, or collected from a pharmacy

See further information that can be given to visitors detailing the practicalities of conducting testing at home (<https://www.gov.uk/government/publications/coronavirus-covid-19-testing-for-adult-social-care-settings>) and registering results to the supported living's UQN. Each pack of tests will come with instructions. Visitors are not able to test themselves at home with a test from a pack of 25 in line with MHRA regulations.

When considering the most appropriate testing route, managers should consider any additional risks that may arise from testing off site, as well as the confidence and ability of visitors to carry out tests away from the care home. This may include factors such as:

- visitors inaccurately conducting or reporting lateral flow testing themselves
- the increased risk of visitors needing to take public transport to a testing site, particularly where it is far from the setting, or coming into contact with other people
- visitors may not have a mobile phone or email address to receive the result of their test

Where visitors will be self-testing, managers may wish to supervise the first few tests on site and provide support to ensure visitors are confident conducting the tests at home and they are being completed and reported satisfactorily.

Wherever the test is conducted, it should be done on the day of the visit. Once the visitor has registered the test, they will receive confirmation of their result by text message (SMS) and email to show proof of result. Proof of a negative result may include an email or text from NHS Test and Trace or a date-stamped photo of the test cartridge itself. Managers do not need to retain records of proof. All tests done at home should be registered to the UQN of the supported living setting and managers should ensure visitors are aware of their UQN and the legal duty to report the result. Where individuals are testing for multiple purposes (for example, if they're part of a school bubble), the test only needs to be linked to one organisation. Being able to link visitors to a supported living setting enables public health teams to better support settings to reduce the transmission of coronavirus and prevent outbreaks.

If the visitor tests positive they and their household must immediately self-isolate, following government guidance for households with possible or confirmed coronavirus (COVID-19) infection (<https://www.gov.uk/government/publications/covid-19-stay-at-home-guidance>). If the test has been taken away from their own home, when returning home, they should avoid public transport and wear a mask. Visitors must also complete a confirmatory polymerase chain reaction (PCR) test, which should be provided to them by the setting if testing on site, or ordered from the government portal (<https://www.gov.uk/get-coronavirus-test>) or by calling 119. This can be returned either through a courier or through a Royal Mail priority postbox. If the confirmatory PCR comes back positive, their household must also self-isolate and contacts may also need to self-isolate.

Visitors who have recently tested positive for COVID-19 with a PCR test should not be retested within 90 days unless they develop new symptoms or unless specific infection detection and response plans are in place for individuals or in the local area already. This means that some visitors will not need to be tested regularly because they will fall into this 90-day window. These visitors should use the result of their positive PCR result to show that they are currently exempt from testing until the 90-day period is over following their period of self-isolation. Once the 90-day period is over, visitors can then continue to be tested. They should still continue to follow all other relevant JPC measures throughout these 90 days, including social distancing, maintaining good hand hygiene and wearing PPE.

For visits taking place at the setting, the manager may also wish to consider:

- if a setting has a communal garden area which can be accessed without anyone going through a shared building, then using this space for visits should be encouraged, if social distancing measures are met
- if in shared accommodation, visitors should avoid (or minimise if avoidance is not possible) contact with other people who live there and staff (with face-to-face contact occurring for less than 15 minutes and at least 2 metres apart). Where needed, conversations with staff can be arranged over the phone following an in-person visit
- visitors should be reminded and provided facilities to wash their hands for 20 seconds or use hand sanitiser on entering and leaving the home, and to catch coughs and sneezes in tissues and clean their hands after disposal of the tissues
- PPE should be viewed as a tool to mitigate risk, rather than a requirement for people to visit settings. Managers may wish to provide visitors with appropriate PPE where visiting is judged to be high-risk and the PPE can be tolerated by both the visitor and the person being visited. If PPE is provided, it should be used in accordance with the guidance and visitors must be reminded that it is only effective if used correctly and combined with infection prevention and control measures such as hand hygiene and avoiding touching your face with your hands. Staff should provide visitors with guidance on how to safely put on and remove PPE and visitors should also be encouraged to view the video demonstration. Visitors should be informed about local arrangements for the disposal of PPE. If managers wish to provide visitors with PPE, this is available to them for free and they can obtain this from the same route as their usual COVID-19 PPE.

For visits taking place away from the setting, the manager should consider:

- offering support so people can find or go to outside spaces to see their relative in a safer environment in line with current social distancing rules
- factors to minimise the risk for staff and other individuals in the supported living setting (including the layout of the premises and the nature of the support provided)
- the nature and context of the visit – for example, whether the visit would include overnight stays in the family home or visits to a public place
- the support needs that the person may have during the visit, and whether they will need to be accompanied by a staff member, carer, family member or friend
- transport for the visit should avoid exposing the person to those outside the household they are visiting, for instance by travelling in a family car wherever possible
- increased communal risks that may arise in shared areas when people return from off-site visits (including shared spaces indoors and outdoors, on-site grouped services and social activities)
- the need for those returning from off-site visits to self-isolate if they develop symptoms, test positive for coronavirus, or have been identified as being in contact with someone who has tested positive for COVID-19. There is no expectation for someone to self-isolate for 14 days after a visit has happened if this is not the case

And in all cases, the manager should work with people being supported and their families to:

- make sure that no one with COVID-19 symptoms should participate in a visit and anyone with suspected symptoms should be tested
- make sure that no one visits who should be self-isolating as they have been a close contact of a COVID-19 case in the previous 10 days, nor anyone who has returned from certain countries (<https://www.gov.uk/guidance/how-to-quarantine-when-you-arrive-in-england>) in the same time period

- those involved in the visit will be supported to follow good infection control practice including social distancing, hand hygiene and face coverings; and whether their needs are likely to impact their ability to do so
- encourage visitors to wear appropriate face coverings or use any other appropriate PPE as advised when visiting to protect people
- in some circumstances, subject to risk assessment, visors may be preferable to masks, to facilitate the more effective provision of care and social interaction through non-verbal communication, but social distancing should be maintained wherever possible
- where possible, visitors can be given support on how to prepare for a visit and given tips on how to communicate if face coverings are required

## Testing of people being supported to assist visiting

Setting managers may decide to use some of their additional rapid lateral flow testing allocation for testing of the person being supported (<https://www.gov.uk/government/publications/coronavirus-covid-19-testing-service-for-extra-care-and-supported-living-settings>). This can be used to facilitate safer activities by testing the person being supported with rapid lateral flow tests. The use of tests is at the managers discretion and must be conducted with the consent of the person being supported. This should be in addition to the regular PCR testing for all people in high-risk supported living and extra care settings.

In cases where a person living in supported living receives a positive result using rapid lateral flow testing or has symptoms, they should self-isolate immediately and take a confirmatory PCR test from the stocks of the setting or get a free test (<https://www.gov.uk/get-coronavirus-test>). If they have been identified as a contact of a confirmed COVID-19 case, they should isolate and follow government guidance (<https://www.gov.uk/government/publications/guidance-for-contacts-of-people-with-possible-or-confirmed-coronavirus-covid-19-infection-who-do-not-live-with-the-person>). Please also alert your local health protection team (HPT) and consider any outbreak measures you have in place.

Rapid lateral flow testing should not be seen as a condition of people in supported living taking part in visits in and out of the setting, but as a tool to support safer visits alongside other JIC measures. For clarity people who test positive or those who are known contacts of a positive case should be isolated according to public health guidance (<https://www.nhs.uk/conditions/coronavirus-covid-19/self-isolation-and-treatment/when-to-self-isolate-and-what-to-do/>), and testing cannot be used during an isolation period to enable earlier release.

Managers may wish to consider testing people living in high risk settings with rapid lateral flow tests if they are often leaving the premises to meet or visit people. This is similar to testing for people who are unable to work from home who can access twice weekly lateral flow testing from their local asymptomatic testing site. Twice weekly testing for people who live in supported living can be conducted on site (<https://www.gov.uk/government/publications/coronavirus-covid-19-lateral-flow-testing-in-adult-social-care-settings>), assisted by a staff member.

## If a supported living worker has COVID-19 symptoms

General guidance for working safely during coronavirus (<https://www.gov.uk/guidance/working-safely-during-coronavirus-covid-19>) is available. If a supported living worker develops COVID-19 symptoms (<https://www.nhs.uk/conditions/coronavirus-covid-19/self-isolation-and-treatment/when-to-self-isolate-and-what-to-do/>), then NHS advice (<https://www.nhs.uk/conditions/coronavirus-covid-19/>) is available. In addition:

- if symptoms start at home (off-duty), they should not attend work and should notify their line manager immediately
- if symptoms start at work, and the staff member is not wearing a face mask they should immediately put on a surgical face mask, inform their line manager and return home
- if symptoms start at work, and the staff member is wearing a face mask, then this and any other PPE should be removed and disposed of carefully, as described in the domiciliary care resource ([https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/892496/Domiciliary\\_guidance\\_v2\\_15Jun.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/892496/Domiciliary_guidance_v2_15Jun.pdf)), hands must be washed, the staff member should immediately put on a surgical face mask, inform their line manager and return home
- staff should get tested as soon as possible and be asked to support all public health requests from NHS Test and Trace (<https://www.gov.uk/guidance/nhs-test-and-trace-how-it-works>) and the local health protection teams
- if symptoms do not improve after 7 days, or their condition gets worse, they should speak to their occupational health department if they have one or use the NHS 111 online (<https://111.nhs.uk/>) coronavirus service. If they do not have internet access, they should call NHS 111. For a medical emergency, they should call 999

More details are available from the stay at home guidance (<https://www.gov.uk/government/publications/covid-19-stay-at-home-guidance>) and the management of exposed staff and patients in health and social care settings (<https://www.gov.uk/government/publications/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings>). Staff should not attend work, visit or care for individuals until safe to do so, as described in the staff return to work criteria ([https://www.gov.uk/government/publications/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings#staff-return-to-work-criteria](https://www.gov.uk/government/publications/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings#staff-return-to-work-criteria)).

Currently it is not known how long any immunity to COVID-19 might last. If a staff member becomes unwell again, they should self-isolate and may need to be tested again.

## Test and Trace

NHS Test and Trace (<https://www.gov.uk/guidance/nhs-test-and-trace-how-it-works>) has been established to help identify, contain and minimise the transmission of COVID-19. This will help to reduce the spread of the virus and save lives. The service is designed to:

- provide testing for anyone who has symptoms of COVID-19 to find out if they have the virus
- get in touch with anyone who has tested positive for COVID-19 to help them share information about any close, recent contacts that they have had
- notify those contacts, where necessary, with instructions to stay at home and self-isolate to help stop the spread of the virus

For health and social care worker contacts, NHS Test and Trace will consider whether full medical-grade PPE has been worn in accordance with current guidance on infection prevention and control (<https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control>). This is to assess whether a health or social care worker should be classed as a close contact and asked to self-isolate.

If you are judged by NHS Test and Trace to have had relevant, close contact with someone who has COVID-19, you must stay at home and self-isolate immediately for 10 days and follow the health advice that you will be directed to. For people in supported living arrangements, this 10-day period should increase to 14 days based on a risk assessment if the setting is considered high risk.

## **If a supported living worker is concerned they may have been exposed to COVID-19**

If a worker or volunteer has come into close contact with a person who is confirmed or suspected of having COVID-19 while not wearing PPE, or had a breach in their PPE, whether within or outside the work setting, then the staff member should inform their line manager, and follow guidance for the management of exposed healthcare workers (<https://www.gov.uk/government/publications/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings>).

For more information on interpreting test results and the actions required for both symptomatic and asymptomatic individuals, see the flowcharts illustrating the return to work process (<https://www.gov.uk/government/publications/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings>).

## **If a supported living worker has symptoms or tests positive (even when asymptomatic)**

If a supported living worker is symptomatic, they must stay at home, self-isolate immediately and order a test. Members of their household must self-isolate too. If the test result is negative and the supported living worker does not have COVID-19 and is then well, they can end their period of isolation. If the worker is still ill, though not with COVID-19, for example they have the flu, they should not return to work. Other members of the household do not have to stay at home and isolate if the test result for COVID-19 is negative.

If a supported living worker tests positive for COVID-19, they must self-isolate for 10 days from the date of the test, even if they are asymptomatic. If they remain asymptomatic, they can return to work on day 11. If, during the 10-day isolation, the supported living worker subsequently develops symptoms, they must self-isolate for 10 days from the day the symptoms started.

On testing positive, the supported living worker will be contacted by NHS Test and Trace and directed to a website to input the details of their close, recent contacts. If the worker is unable to use a web-based system, they will receive a phone call from a health professional. As a supported living worker is employed in a health and care setting, the contact tracing process will be escalated to local public health experts, who will liaise with the manager of the relevant setting, if necessary.

The household of the supported living worker should stay at home and self-isolate for 10 days from the day the test was taken. If any member of the household then develops symptoms of COVID-19 during the 10-day period, they should continue to stay at home and isolate for 10 days after the onset of their symptoms, in line with the stay at home guidance (<https://www.gov.uk/government/publications/covid-19-stay-at-home-guidance/stay-at-home-guidance-for-households-with-possible-coronavirus-covid-19-infection>). The individual should also arrange a test to check if they have COVID-19.

If a supported living worker is tested while asymptomatic and has a negative result for COVID-19, they can return to work.



## If someone in supported living has symptoms of COVID-19

If the person develops a COVID-19 infection, plans need to be developed so that the person is supported to have their health checked in case additional help and health interventions are needed.

It may be harder to recognise COVID-19 infection in people with dementia, autistic people, and people with learning disabilities who may not be able to communicate verbally or easily express the symptoms they are experiencing. This includes signs of a high temperature (37.8°C or above), a cough, or a change in sense of taste or smell, as well as for softer signs, ie being short of breath, being not as alert, having a new onset of confusion, being off food, having reduced fluid intake, diarrhoea or vomiting.

Annex A sets out special considerations when taking swab samples from people who may find the process challenging. It is essential that processes are put in place to enable and ensure the healthcare of these people is effectively supported.

This should include delivery of a rights-based approach such as:

- consideration of all possible diagnostic causes
- access to healthcare
- informing individuals of their rights and ability to challenge decisions
- access advocacy
- use of the hospital passport
- regularly consulting with family members and carers

If a person being supported develops symptoms, then these plans should be acted on to provide additional support and help them self-isolate, and ensure that visitors such as care/support workers and family members follow appropriate procedures such as handwashing, respiratory hygiene and, where appropriate, the use of P.P.E.

It will be important to be aware of the specific needs of people living with dementia in this regard. For example, people with dementia may not fully understand the significance of, and need for, isolation. They may also find it frightening to see a carer wearing P.P.E. Simple actions can be taken to mitigate against this such as having the supported care worker's name and picture clearly visible on clothing, using tone of voice and open body language to demonstrate warmth and drawing or using written words to communicate where appropriate

Plans should include communications to family and others who provide support to help understand the reasons for staying in isolation. The principles underpinning the Mental Capacity Act (<https://www.scie.org.uk/mca/introduction/mental-capacity-act-2005-at-a-glance>) (2005) should be followed when it is felt a person being supported may lack capacity to make a decision.

Autistic people, people with mental ill health, learning disabilities and dementia may need support in keeping isolated from the people with whom they may share communal facilities. If isolation is not possible within the supported living service, then appropriate alternative community provision may need to be considered. This needs to be discussed with the individual and, where appropriate, family members.

## Managing outbreaks in supported living settings

The Health Protection Teams (HPTs) at PHE have an essential role in responding to and supporting any infectious disease outbreaks in supported living settings. Your local HPT will provide tailored infection prevention control advice to ensure staff protect themselves and the people they support.

To confirm the presence of an outbreak, PHE is responsible for the initial risk assessment and initiating testing of suspected outbreaks in supported living settings (depending on local systems in place with other stakeholders such as local authorities and the NHS).

If an outbreak is suspected in a supported living setting, this should be reported to the local HPT immediately. They will undertake an initial risk assessment, provide advice on outbreak management, and decide what testing is needed. Local HPTs will also inform their local partners of the situation.

Find details for your local HPT and more information (<http://www.gov.uk/health-protection-team>).

With consent, the person's GP should be informed if the person who uses services has signs and symptoms compatible with COVID-19.

An outbreak in, or associated with, a supported living setting is defined as within a 14-day period:

1. there are 2 or more confirmed or suspected cases of COVID-19 in a supported living environment
2. a care worker becomes aware that more than one person they support has COVID-19 symptoms, or
3. a care worker and a person who receives care from this worker have COVID-19 symptoms

During an outbreak or when an outbreak is suspected:

- symptomatic people should self-isolate
- if there are any communal areas in the setting which cannot be avoided, then people who are symptomatic or have tested positive for COVID-19 should not attend these communal areas at the same time as others and these areas should be cleaned after use
- if bathrooms or lavatories are not available for sole use by an individual who has tested positive, strict cleaning protocols must be implemented in shared bathroom or lavatory facilities after each use. Where appropriate, any people being supported who are asymptomatic should use separate facilities to those who are symptomatic. The landlord or the care provider should provide a deep cleaning function to all facilities to enable consistent good hygiene practice. For more information on deep cleaning, please refer to the COVID-19 deep cleaning in care homes guidance (<https://www.infectionpreventioncontrol.co.uk/content/uploads/2020/05/COVID-19-Deep-cleaning-guidance-for-Care-Homes-May-2020.pdf>)
- where the supported living environment is cleaned by the tenant, advice and guidance should be offered

## Testing

### COVID-19 testing for supported living staff and people being supported

All supported living staff displaying COVID-19 symptoms (including symptomatic members of their household) can access a test, as confirmed in the government's adult social care action plan (<https://www.gov.uk/government/publications/coronavirus-covid-19-adult-social-care-action-plan>).

If you are a member of staff and need a COVID-19 test because you are symptomatic, you should self-isolate for at least 10 days from when symptoms started and get tested (<https://www.gov.uk/get-coronavirus-test>).

Anyone experiencing coronavirus symptoms can now be tested, which includes people receiving care and support. This can be accessed through the digital portal (<https://www.nhs.uk/conditions/coronavirus-covid-19/testing-and-tracing/ask-for-a-test-to-check-if-you-have-coronavirus/>) or via NHS 111 service to book testing.

For people who lack the capacity to consent to testing for themselves, the Mental Capacity Act issues described above apply.

### **Testing for patients and discharge from hospital into the community**

Some people with non-urgent needs, who do not meet the clinical criteria to reside in hospital, will be discharged for their recovery period.

As set out in the COVID-19 adult social care action plan (<https://www.gov.uk/government/publications/coronavirus-covid-19-adult-social-care-action-plan/covid-19-our-action-plan-for-adult-social-care>), any individual moving into a supported living setting should be supported as if they were possibly COVID-19-positive until a 10-day period has passed, even where they have tested negative for COVID-19. This should increase to 14 days based on a risk assessment if the setting is considered high risk. Providers will need to follow the relevant guidance for use of PPE (<https://www.gov.uk/government/publications/covid-19-how-to-work-safely-in-domiciliary-care>) for COVID-19-positive people during this period.

All people receiving hospital care will be tested for COVID-19, and hospitals should share care needs and COVID-19 status with relevant community partners planning the subsequent community care. Supported living environments should ensure that support plans are in place to maintain a supportive and planned transfer and are discussed with the person being discharged, and where appropriate their family and care providers.

If the PCR (swab) test has been performed in hospital but the result still awaited, the person may only be discharged if assurance has been gained that appropriate support plans are in place for the requirements of the isolation period to be met.

For autistic people and people with learning disabilities, mental ill health, or dementia it will be particularly important to make sure they and their families understand, before the transfer happens:

- why these arrangements are needed
- what they will look like
- how long they will go on for

It will be important to ask how they feel about this and what, if anything, could make it easier for them.

For people living in shared settings, the views and needs of the people they live with should be taken into account when thinking through the practical arrangements that need to be made at home to reduce the risk of infection, such as that set out in the guidance for households with possible or confirmed coronavirus (COVID-19) infection (<https://www.gov.uk/government/publications/covid-19-stay-at-home-guidance/stay-at-home-guidance-for-households-with-possible-coronavirus-covid-19-infection#why-staying-at-home-is-very-important>). This should include their view on the risk to them of sharing their home with the person

being discharged whose COVID-19 status is unknown and may include the need to consider if any change in living arrangements is needed, for example to allow the person being discharged to access a separate bathroom and stopping other people visiting the home.

Some living in the home may need support to make sure that advice about what should happen during the isolation period can be followed. This could include not having visitors to the house and cleaning shared areas such as kitchens after use.

If the person being discharged, or anyone they share their home with, lacks capacity to understand information about the discharge arrangements or the requirements of the isolation period, and decisions need to be made that impact on their living arrangements and/or support needs, then the Mental Capacity Act should be followed and people who are significant to them consulted with before decisions are taken in their best interests. Prior to this happening, all steps should be taken to support the person to understand information, which may include using accessible formats such as easy read or having the support of someone who knows them well to communicate information.

## Personal protective equipment (PPE)

The risk of transmission should be minimised through safe working procedures, reducing contact and following standard JPC precautions as described in the how to work safely in domiciliary care (<https://www.gov.uk/government/publications/covid-19-how-to-work-safely-in-domiciliary-care>) resource.

The information on PPE below references supported living staff, but also applies to other care providers coming into the environment to provide care. The supported living manager should work with the people who live there and their families to ensure that external care providers follow the guidance. Updated guidance on PPE (<https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control/covid-19-personal-protective-equipment-ppe>) should be referred to if aerosol generating procedures are being carried out.

Please refer to the correct order of donning

([https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/882069/Putting\\_on\\_PPE\\_home\\_carer.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/882069/Putting_on_PPE_home_carer.pdf)) (putting PPE on) and doffing

([https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/882070/Taking\\_off\\_PPE\\_home\\_carer.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/882070/Taking_off_PPE_home_carer.pdf)) (taking PPE off) PPE. For AGPs (<https://www.gov.uk/government/publications/covid-19-personal-protective-equipment-use-for-aerosol-generating-procedures>) (aerosol generating procedures) please follow specific PPE guidance. PPE should always be used in accordance with JPC measures and requirements for hand hygiene should include washing of exposed forearms.

Table 1: when providing close personal care in direct contact with the person(s) in a supported living setting (for example, touching) or within 2 metres of anyone in the setting who is coughing

Recommended PPE items	Explanation
Disposable gloves	Single use to protect you from contact with the person's body fluids and secretions.
Disposable plastic apron	Single use to protect you from contact with the person's body fluids and secretions.

Recommended PPE items	Explanation
Fluid-repellent (Type IIR) surgical mask	<p>Fluid-repellent surgical masks (FRSMs) can be used continuously while providing care, unless you need to remove the mask from your face (for example to drink, eat, take a break from duties). You can wear the same mask between different home care visits (or visiting different people living in an extra care scheme), if it is safe to do so whilst travelling. This may be appropriate when travelling between households on foot or by car or by public transport, so long as you do not need to take the mask off, or lower it from your face and providing it does not compromise your safety (for example, driving ability) in any way. You should not touch your face mask. The mask is worn to protect you, the care worker, and can be used while caring for a number of different people regardless of their symptoms. You should remove and dispose of the mask if it becomes damaged, visibly soiled, damp, or uncomfortable to use. If removed, you would then need to use a new mask when you start your next home care visit. Technical Specifications for <u>PPE</u> (<a href="https://www.gov.uk/government/publications/technical-specifications-for-personal-protective-equipment-ppe">https://www.gov.uk/government/publications/technical-specifications-for-personal-protective-equipment-ppe</a>)</p>
Eye protection	<p>Eye protection is recommended for care of people where there is risk of droplets or secretions from the person's mouth, nose, lungs or from body fluids reaching the eyes (for example, caring for someone who is repeatedly coughing). Use of eye protection should be discussed with your manager and you should have access to eye protection (such as goggles or visors). Eye protection can be used continuously while providing care, unless you need to remove the eye protection from your face (for example, to take a break from duties). We do not recommend continued use of eye protection when driving or cycling. If you are provided with goggles/a visor that is reusable, then you should be given instructions on how to clean and disinfect following the manufacturer's instructions or local infection control policy and store them between visits. If eye protection is labelled as for single use then it should be disposed of after removal.</p>

The recommendations in Table 2 below apply when within 2 metres of a person but not delivering personal care or needing to touch them, and there is no one within 2 metres who has a cough.

This includes:

- for tasks such as: removing medicines from their packaging, prompting people to take their medicines, preparing food for people who can feed themselves without assistance, or cleaning
- whatever your role in care (ie applies to all staff, care workers, cleaners etc). If practical, household members with respiratory symptoms should remain outside the room or rooms where the care worker is working. They should be encouraged to follow good hand and respiratory hygiene and remain 2 metres away
- if unable to maintain 2-metre distance from anyone in the household who is coughing (including the person receiving care/support) then follow recommendations in Table 1 above

These principles are also suitable for extra care housing schemes. It is important to note that PPE is only effective when combined with:

- hand hygiene (cleaning your hands regularly and appropriately)
- respiratory hygiene and avoiding touching your face with your hands
- following standard infection prevention and control precautions:
  - Healthcare-associated infections: prevention and control in primary and community care (<https://www.nice.org.uk/guidance/cg139>) (NICE)
  - Standard infection control precautions: national hand hygiene and personal protective equipment policy ([https://improvement.nhs.uk/documents/4957/National\\_policy\\_on\\_hand\\_hygiene\\_and\\_PPE\\_2.pdf](https://improvement.nhs.uk/documents/4957/National_policy_on_hand_hygiene_and_PPE_2.pdf)) (NHS England and NHS Improvement)

Table 2: when within 2 metres of a person but not delivering personal care or needing to touch the person(s) in a supported living setting, and there is no one within 2 metres who has a cough

<b>Recommended PPE item</b>	<b>Explanation</b>
Disposable gloves – sometimes required*	* Required if for other reasons set out in standard infection prevention and control precautions (for example, contact with person’s bodily fluids) or if anyone in the household is shielding.
Disposable plastic apron – sometimes required*	* Required if for other reasons set out in standard infection prevention and control precautions (for example contact with person’s bodily fluids) or if anyone in the household is shielding.

Recommended PPE item	Explanation
Type II surgical mask – required	<p>Type II surgical masks can be used continuously while providing care, unless you need to remove the mask from your face (for example, to drink, eat, take a break from duties). You can wear the same mask between different home care visits (or visiting different people living in an extra care scheme), if it's safe to do so while travelling. This may be appropriate when travelling between households on foot or by car or by public transport, so long as you do not need to take the mask off, or lower it from your face and providing it does not compromise your safety (for example, driving ability) in any way.</p> <p>You should not touch your face mask.</p> <p>The mask can be used while caring for a number of different people regardless of their symptoms. You should remove and dispose of the mask if it becomes damaged, visibly soiled, damp, or uncomfortable to use. If removed, you would then need to use a new mask when you start your next home care visit.</p> <p>Note: surgical masks do not need to be fluid repellent for use in this situation. However, if you are already wearing a fluid-repellent (Type IIR) surgical mask there is no need to replace it, and if only fluid-repellent (Type IIR) surgical masks are available then these may be used.</p> <p>If the next visit you undertake includes personal care, then you will need to follow recommendations in Table 1 for the next visit. Details on specification can be found (<a href="https://www.gov.uk/government/publications/technical-specifications-for-personal-protective-equipment-ppe">https://www.gov.uk/government/publications/technical-specifications-for-personal-protective-equipment-ppe</a>)</p>
Eye protection	Not required

## Cleaning

In supported living environments, cleaning may be carried out by the person who lives there, their family, an external cleaner, or as a service provided as part of the accommodation. Where appropriate the supported living manager should adapt guidance accordingly. Additional guidance is available for cleaning in non-healthcare settings (<https://www.gov.uk/government/publications/covid-19-decontamination-in-non-healthcare-settings/covid-19-decontamination-in-non-healthcare-settings>).

If the person you are supporting does their own cleaning or arranges their own cleaner, then staff should consider how the person they are supporting is most likely to understand the information and use the most appropriate communication techniques for that person. With their consent it may be appropriate to place visible guidance such as pictorial posters, or other communication aides to reinforce cleaning messaging.

If the person in the supported living setting has COVID-19 symptoms or has confirmed COVID-19, then their personal waste (for example, used tissues, continence pads and other items soiled with bodily fluids) and disposable cleaning cloths can be discarded in clinical waste bins, where available, or can be stored securely within disposable rubbish bags. These bags should be placed into another bag, tied

securely and kept separate from other waste which can be disposed of as per usual practice. This should be put aside for at least 72 hours before being put in the usual household waste bin for disposal as normal. Where the person has a learning disability or other needs, it will be important to make sure they understand exactly what they need to do and why.

## Laundry

In supported living environments, laundry may be carried out normally by the person who lives there, their family, an external person, or as a service provided as part of the accommodation. Where appropriate the supported living manager should adapt guidance accordingly. If a laundry service is provided, it should follow the guidance below:

- wash items in accordance with the manufacturer's instructions. Use the warmest water setting and dry items completely. Dirty laundry that has been in contact with an unwell person can be washed with other people's items
- do not shake dirty laundry; prior to washing this minimises the possibility of dispersing the virus through the air
- clean and disinfect anything used for transporting laundry with your usual products, in line with the cleaning guidance above

If someone carries out their own laundry duties then with consent it may be appropriate to place visible pictorial reminders, such as posters, or other communication aides in line with the persons individual communication method around the supported living setting, to reinforce the above laundry messaging.

## Annex A: taking swabs

This annex sets out special considerations when taking samples from people who may find it difficult to understand what is happening. This could include autistic people and people with learning disabilities, mental ill health, dementia or any other type of cognitive impairment:

- be aware that the person has a cognitive impairment that may impact on their ability to understand information about taking the swab
- find out from those who know them best how and when to give the person information about taking the swab in a way they are most likely to understand it
- having given the information, if it is concluded that the person does not have the mental capacity to understand it and consent to taking the swab, a decision should be made in their best interests following the principles of the Mental Capacity Act 2005. See Coronavirus (COVID-19): looking after people who lack mental capacity (<https://www.gov.uk/government/publications/coronavirus-covid-19-looking-after-people-who-lack-mental-capacity>)
- relevant information about the person's needs, preferences and understanding should be taken into account, and where possible, a family member or carer who knows them well should be present or at least consulted with to inform the best interests decision
- provide reassurance and use a calm and confident approach
- explain the process step by step using appropriate language and their preferred communication methods. If appropriate, use visual aids to show what is happening



- be prepared to take time when taking the sample and to try more than once if needed, possibly at different times of the day
- if the person becomes distressed at any point, it may be necessary to abandon the attempt to take a sample

## Annex B: additional resources

- The Social Care Institute for Excellence has produced guidance for care staff who support autistic people and people with learning disabilities (<http://www.scie.org.uk/care-providers/coronavirus-covid-19/learning-disabilities-autism>)
- The Alzheimer's Society website has resources to promote awareness of the Herbert Protocol among local emergency services and the local community. The Herbert Protocol is a national scheme that encourages carers, family and friends to provide and put together useful information, which can be used in the event of a vulnerable person going missing (<https://www.alzheimers.org.uk/get-support/publications-and-factsheets/dementia-together-magazine/scheme-support-missing-people>)
- Examples of factsheets developed by organisations include 'This is Me' (<http://www.alzheimers.org.uk/get-support/publications-factsheets/this-is-me>), which contains space for:
  - important routines
  - access to advanced care plans
  - cultural, spiritual, religious and family background
- The Challenging Behaviour Foundation (<https://www.challengingbehaviour.org.uk/information/covid19information.html#Infosheets>) has produced an information sheet about people with severe learning disabilities and face masks. The resource provides useful information about helping people with severe learning disabilities to prepare for the experience of wearing or seeing other people wear P.P.E.
- Organisations such as Speakup (<http://www.speakup.org.uk/coronavirus>) have developed a series of resources such as hospital passports for autistic people or people with a learning disability to help on discharge to hospital, if this was necessary.
- The Housing Learning and Improvement Network have produced various resources related to specialist housing and COVID-19. These include bereavement and emotional support materials (<http://www.housinglin.org.uk/Topics/type/Coronavirus-COVID-19-Top-tips-in-Bereavement-Care-in-Specialist-Housing/>).

Print this page