




Royal College of  
Obstetricians &  
Gynaecologists

# Coronavirus (COVID-19) Infection in Pregnancy

---

Information for healthcare professionals

Version 10.1: Published Friday 19 June 2020



# Contents

	Summary of updates	3
<b>1</b>	Purpose and scope	4-9
<b>2</b>	Antenatal care during the COVID-19 pandemic	10-16
<b>3</b>	Venous thromboembolism prevention	17-19
<b>4</b>	Labour and birth during the COVID-19 pandemic	20-27
<b>5</b>	Managing clinical deterioration during the COVID-19 pandemic	28-33
<b>6</b>	Postnatal care	34-37
<b>7</b>	Acknowledgments	38
<b>8</b>	Appendix 1: Summary of previous changes	39-48
<b>9</b>	Appendix 2: Key considerations when caring for symptomatic women with suspected/confirmed COVID-19	49-52
<b>10</b>	References	53-60

## Summary of updates

Previous updates have been summarised in Appendix 2. New updates for this version of the guideline are summarised here.

Version	Date	Summary of changes
<b>10.1</b>	19.6.20	<b>1.1:</b> Removal of 'MERS, Middle East Respiratory Syndrome' from the literature search strategy since it has not resulted in any new references since the first search.
<b>10.1</b>	19.6.20	<b>1.4:</b> UKOSS reference changed to the published article in <i>The BMJ</i> .
<b>10.1</b>	19.6.20	<b>2.2:</b> Advice on face masks changed to reflect national guidance from NHS England.
<b>10.1</b>	19.6.20	<b>4.4:</b> Advice on number of visitors and/or birth partners for hospital inpatients changed to reflect national guidance from NHS England.
<b>10.1</b>	19.6.20	<b>5.2:</b> Advice for women who are clinically deteriorating modified to include government recommendations based on the interim results of the RECOVERY trial.
<b>10.1</b>	19.6.20	<b>6.2:</b> Specified that babies should not be advised to wear face masks because of the risk of suffocation.



Royal College of  
Obstetricians &  
Gynaecologists

# I. Purpose and scope

# I. Purpose and scope

This document is designed to provide guidance to healthcare professionals who care for pregnant women during the COVID-19 pandemic. It is designed to provide advice on how existing clinical guidelines can be implemented during this time, not to replace them.

The advice in this document is provided as a resource for UK healthcare professionals based on a combination of available evidence, good practice and expert consensus opinion. The priorities are:

- (i) The reduction of transmission of COVID-19 to pregnant women.
- (ii) The provision of safe, personalised and woman-centred care during pregnancy, birth and the early postnatal period during the COVID-19 pandemic.
- (iii) The provision of safe, personalised and woman-centred care to pregnant and postnatal women with suspected/confirmed COVID-19.

Please be aware that this is very much an evolving situation and this guidance is a living document that is being updated as new information becomes available. We therefore suggest that you visit this page regularly for current advice. This guidance will be kept under regular review as new evidence emerges. If you would like to suggest additional areas for this guidance to cover, any clarifications required or to submit new evidence for consideration, please email [COVID-19@rcog.org.uk](mailto:COVID-19@rcog.org.uk). Please note, we will not be able to give individual clinical advice or information for specific organisational requirements via this email address.

[Information for pregnant women and their families](#) is available in question and answer format, with accompanying videos in some cases, on the RCOG COVID-19 hub.

## I.1 Identification and assessment of evidence

This guidance document is updated regularly following a review of the evolving literature during this pandemic. Weekly literature reviews are generated using the following search terms, MESH headings and associated synonyms: pregnancy, coronavirus, SARS, severe acute respiratory syndrome, infant, newborn and breast feeding. The search results are published weekly on the [RCOG website](#).<sup>1</sup>

Due to the duration and rapidly evolving nature of the COVID-19 pandemic, there is a current lack of high-quality evidence. Using a conventional grading system for guideline development, such as SIGN, many of the studies would be classed as level 3 or 4 (non-analytical studies, e.g. case series/reports and expert opinion), with a few recent studies being classed as level 2 (systematic reviews of cohort studies).<sup>2</sup> The advice based on this evidence would therefore be graded D, and in some cases, graded as good practice points. Good practice points were defined based on expert consensus opinions of senior clinicians across a variety of disciplines reviewing the available data from case series, ongoing studies and clinical practice.

As this is an evolving pandemic, this guidance is updated regularly. Most advice is extrapolated from case reports or systematic reviews of case series, along with expert consensus opinion to develop the advice statements. Clinicians and patients are advised to be aware of the low-quality evidence on which the advice is given when using this guidance to assist decision making.

## 1.2 Epidemiology

Novel coronavirus (SARS-CoV-2) is a new strain of coronavirus causing COVID-19, first identified in Wuhan City, China towards the end of 2019.<sup>3</sup> Other human coronavirus (HCoV) infections include HCoV 229E, NL63, OC43 and HKU1, which usually cause mild to moderate upper-respiratory tract illnesses, like the common cold, Middle East Respiratory Syndrome (MERS-CoV) and Severe Acute Respiratory Syndrome (SARS-CoV).<sup>4</sup>

The World Health Organization (WHO) publishes a daily [international situation report](#) with an additional [Situation Dashboard](#) to provide information for individual countries. The [total number of confirmed cases](#) in the UK is published by the Department of Health and Social Care, and is available in a [visual dashboard](#).

For the most up-to-date advice please refer to health protection agency websites for [England](#), [Wales](#), [Scotland](#) and [Northern Ireland](#). Public Health England (PHE) and Public Health Scotland (PHS) have been cited throughout this document; specific guidance from the other areas of the UK will be updated as they become available. At the time of writing, Public Health Wales are aligning with PHE on case definitions, assessment, infection prevention and control (IPC) and testing. We will update the RCOG guidance if these change.

## 1.3 Transmission

Most global cases of COVID-19 have evidence of human-to-human transmission. This virus can be readily isolated from respiratory secretions, faeces and fomites (objects). Transmission of the virus is known to occur through close contact with an infected person (within 2 metres) or from contaminated surfaces.

Pregnant women do not appear more likely to contract the infection than the general population.<sup>5 6</sup> Pregnancy itself alters the body's immune system and response to viral infections in general, which can occasionally cause more severe symptoms. This may be the same for COVID-19 but there is currently no evidence that pregnant women are more likely to be severely unwell needing admission to intensive care or die from the illness than non-pregnant adults.<sup>7</sup>

With regard to vertical transmission (transmission from woman to her baby antenatally or intrapartum), emerging evidence now suggests that vertical transmission is possible.<sup>8</sup> There are, however, serious limitations to the available evidence.<sup>9</sup> Two reports have published evidence of IgM for SARS-CoV-2 in neonatal serum at birth.<sup>10 11</sup> Assuming that IgM does not cross the placenta, this would suggest a neonatal immune response to in utero infection. It is uncertain in these cases whether the IgM levels resulted from cross reactivity as there was no evidence of SARS-CoV-2 in the infant's nasopharyngeal swabs or in the mother's vaginal secretions or breastmilk on PCR testing.<sup>9 12</sup> Moreover, the proportion of pregnancies affected and the significance to a neonate has yet to be determined. In the interim report from the UK Obstetric Surveillance System (UKOSS), 2.5% of babies (n=6) had a positive nasopharyngeal swab within 12 hours of birth.<sup>13</sup> In a systematic review of 24 pregnant women with COVID-19, there was no evidence of SARS-CoV-2 on PCR testing of placenta, amniotic fluid, cord blood or breastmilk samples.<sup>14</sup> Further investigation around vertical transmission is required and is underway.<sup>9</sup>

## 1.4 Effect of COVID-19 on pregnant women

It is known that, whilst pregnant women are not necessarily more susceptible to viral illness, physiological pregnancy related changes to their immune system in pregnancy can be associated with more severe symptoms.<sup>15</sup> This is particularly true in the third trimester.

There is evolving evidence within the general population that there could be a cohort of asymptomatic individuals or those with very minor symptoms who are carrying the virus, although the prevalence is unknown. Most pregnant women will experience only mild or moderate cold/flu-like symptoms. Cough, fever, shortness of breath, headache, anosmia and loss of taste are other relevant symptoms.<sup>16</sup> More severe symptoms which suggest pneumonia and marked hypoxia are widely described with COVID-19 in older people, the immunosuppressed and those with chronic conditions such as diabetes, cancer or chronic lung disease.<sup>17</sup> The symptoms of severe infection are no different in pregnant women and early identification and assessment for prompt supportive treatment is key.

Two case series have been published by clinicians in New York which suggest possible patterns of disease in pregnant women. The first describes 43 pregnant women who tested positive for SARS-CoV-2 and reported a similar pattern of disease severity to non-pregnant adults: 86% mild, 9% severe and 5% critical, although the sample size was too small to draw a definitive conclusion and no comparison was made for age, sex or comorbidity-matched individuals.<sup>18</sup> The second describes the results of screening all 215 women who attended two paired maternity units for labour and birth over a 2-week period.<sup>19</sup> Of these women, 13% (n=33) tested positive for SARS-CoV-2 from nasopharyngeal swabs on attendance to the hospital, although only four (1.9% of all women) had symptoms of COVID-19 on attendance. 88% of those testing positive were asymptomatic.

On Friday 20 March 2020, UKOSS launched a registry for all women admitted to UK hospitals with confirmed COVID-19 in pregnancy. Further information can be [found here](#). An interim report was published on Monday 8 June 2020.<sup>13</sup> The UKOSS study is the largest published population-based cohort of pregnant women admitted to hospital with COVID-19 to date. At the time of the interim report, complete data were available for 427 pregnant women admitted to UK hospitals with confirmed SARS-CoV-2 infection between 1 March and 14 April 2020; this represents an admission rate of 4.9 (95% confidence interval [CI] 4.5–5.4) per 1000 maternities. The women included in the report may have required admission to hospital for many reasons, including severe symptoms of COVID-19 or other obstetric indications (e.g. labour and birth) where COVID-19 was co-existent; it is not known what proportion of the admissions were because of COVID-19, rather than with COVID-19. Of the 427 pregnant women reported in the UKOSS study, 9% required level-3 critical care; four women (<1%) received extracorporeal membrane oxygenation (ECMO).<sup>13</sup> Five women included in the study died, suggesting a SARS-CoV-2-associated maternal mortality rate of 5.6 (95% CI 1.8–13.1) per 100,000 maternities, compared with the maternal mortality rate in the UK of 9.2 per 100,000 in 2015–2017.<sup>20</sup> Whether these deaths are a direct result of COVID-19 infection is currently unclear, and these data are likely to be updated in the future.<sup>13</sup>

In the data from UKOSS, most women were hospitalised in the third trimester or peripartum (n=342, 81%). The median gestational age at hospital admission was 34 completed weeks (interquartile range [IQR] 29–38). Of those admitted, 42% did not require iatrogenic birth of the baby; these women were discharged whilst still pregnant. Of those who did give birth during the data collection period, 59% had caesarean births; approximately half of these were because of maternal or fetal compromise. The remainder were for obstetric reasons (e.g. progress in labour, previous caesarean birth) or maternal request (6%). Of the women having a caesarean birth, 20% required general anaesthesia (GA) because of severe COVID-19 symptoms or urgency of birth.<sup>13</sup>

In the UK, the Intensive Care National Audit and Research Centre (ICNARC) weekly report on individuals admitted to critical care with COVID-19 was updated on 29 May 2020, describing the first 9347 patients admitted to critical care settings with a diagnosis of COVID-19.<sup>7</sup> Of these, 23 were currently pregnant and 34 recently pregnant (within the last 6 weeks). There have also been case

reports of women with severe COVID-19 infection at the time of birth who have required ventilation and ECMO,<sup>21</sup> and of maternal death.<sup>13 22-24</sup> However, the overall numbers are small.

## 1.5 Risk factors for hospital admission with COVID-19 infection in pregnancy

Risk factors that appear to be associated with hospital admission with COVID-19 illness include:

1. Black, Asian or minority ethnicity (BAME)
2. Overweight or obesity
3. Pre-existing comorbidity
4. Maternal age >35 years

The characteristics of women admitted to hospital with COVID-19 in the data from the UKOSS study were compared with controls derived from a historical cohort of women giving birth between 1 November 2017 and 30 October 2018 (n=694). Pregnant women admitted to hospital with COVID-19 during the 2020 pandemic were more likely to be of black or other minority ethnicity (adjusted odds ratio [aOR] 4.49, 95% CI 3.37–6.00), have pre-existing comorbidity (aOR 1.52, 95% CI 1.12–2.06), be aged over 35 years (aOR 1.35, 95% CI 1.01–1.81) or be overweight (BMI of 25–29 kg/m<sup>2</sup>) or obese (BMI 30–39 kg/m<sup>2</sup>; aORs 1.91, 95% CI 1.37–2.68 and 2.20, 95% CI 1.56–3.10, respectively). This suggests that women with these risk factors were disproportionately represented in hospital admissions with or for COVID-19.<sup>13</sup> The association with BAME is particularly apparent and echoes previous findings that UK BAME pregnant women have worse outcomes in maternity.<sup>20</sup> Furthermore, 13% of the UK's total population identifies as being from a BAME background, but 55% of all individuals admitted to UK critical care for COVID-19 illness are from BAME backgrounds and individuals from BAME backgrounds are more likely to die from COVID-19.<sup>7 20 25-27</sup> In the case of COVID-19, it has been postulated that this association may be related to socioeconomic or genetic factors, or differences in response to infection; however, further research is needed.<sup>13 28</sup>

It is estimated that vitamin D deficiency affects over 1 billion people worldwide. Vitamin D deficiency is associated with Acute Respiratory Distress Syndrome (ARDS) which is seen in COVID-19 infection.<sup>29 30</sup> Women of BAME background with melanin pigmented skin often develop vitamin D deficiency; it is estimated that as many as 94% of the South Asian population in the UK are affected by vitamin D deficiency in the winter.<sup>31</sup> Recently, vitamin D supplementation has been suggested to be beneficial in reducing the risk of respiratory tract infections, although data are limited.<sup>32</sup> The current UK advice recommends vitamin D supplementation to all individuals of BAME background, regardless of the COVID-19 pandemic.<sup>33</sup>

In addition to the UKOSS study, which showed that pregnant women with a BMI  $\geq 25$  kg/m<sup>2</sup> were more likely to be admitted to hospital with COVID-19 than the historical controls, other studies in non-pregnant populations have shown a similar trend in terms of worse outcomes for individuals with BMI >25 kg/m<sup>2</sup>.<sup>13</sup> The UK ICNARC weekly report also found that 74% of those patients admitted were overweight or obese; 35% of admitted individuals had a BMI of 25–29 kg/m<sup>2</sup> (overweight), 31% a BMI of 30–39 kg/m<sup>2</sup> (obese) and 7.7% a BMI  $\geq 40$  kg/m<sup>2</sup>.<sup>27</sup>

Pre-existing diabetes mellitus or gestational diabetes affects 5% of pregnant women in the UK, although 88% of women with diabetes in pregnancy are affected by gestational diabetes.<sup>34</sup> In the UKOSS study, comorbidities such as diabetes were associated with pregnant women being admitted to hospital with COVID-19.<sup>13</sup> In non-pregnant individuals, a UK study of 20,133 patients admitted



to high dependency and intensive care with COVID-19 found uncomplicated diabetes was one of the most common comorbidities (21%, 3650/17,599); a further 7% (n=1299) of individuals had complicated diabetes.<sup>35</sup>

Lifestyle measures such as [regular exercise](#), a [healthy diet](#) and [vitamin D supplementation](#) are recommended in pregnancy and throughout life to prevent obesity, type 2 diabetes mellitus and vitamin D deficiency.

## **1.6 Effect of COVID-19 on the fetus**

There are currently no data suggesting an increased risk of miscarriage in relation to COVID-19. Case reports from early pregnancy studies with SARS-CoV and MERS-CoV have not demonstrated a significant relationship between infection and increased risk of miscarriage or second trimester loss.<sup>36</sup>

In the UKOSS cohort, the median gestational age at birth was 38 weeks (IQR 36–39 weeks). Of women who gave birth, 27% had preterm births; 47% of these were iatrogenic for maternal compromise and 15% were iatrogenic for fetal compromise, with 10% of term babies requiring admission to the neonatal unit. Six (2.5%) babies had a positive test for SARS-CoV-2 during the first 12 hours after birth; three of these were in babies born by pre-labour caesarean birth. One of these babies required admission to the neonatal unit. It was unclear from the report whether two perinatal deaths were related to co-existing maternal COVID-19.<sup>13</sup>

A review of 71 neonates delivered to women with COVID-19 in the third trimester reported that neonatal infection was diagnosed in 4 cases (5.6%) within 48 hours of delivery by PCR tests of cord and neonatal blood samples.<sup>8</sup>



Royal College of  
Obstetricians &  
Gynaecologists

## 2. Antenatal care during the COVID-19 pandemic

## 2 Antenatal care during the COVID-19 pandemic

### 2.1 What are the considerations for organisation of antenatal care?

#### Advice

- Women should be advised to continue their routine antenatal care, although it may be modified, unless they meet current self-isolation criteria for individuals or households with suspected or confirmed COVID-19.
- Service modifications are required to assist women practising social distancing measures, to reduce the risk of transmission between women, staff and other clinic/hospital visitors, and to provide care to women who are self-isolating for suspected/confirmed COVID-19 but for whom a hospital attendance is essential.
- Basic assessments such as blood pressure and urine testing are still required. Trusts should plan local strategies to ensure women receive this monitoring.
- Units should employ teleconferencing and videoconferencing where possible and consider which appointments can be most appropriately conducted remotely.
  - The limitations of virtual consultation methods should be recognised, including recognising that some women will not have sufficient internet access on their mobile devices or other computer hardware.
  - It should be acknowledged that virtual appointments, particularly by telephone, may cause new challenges in relationship-building between healthcare professionals and women, especially among vulnerable groups, women for whom English is not their first language or women who are deaf.
  - Healthcare professionals should be aware that women may have additional queries regarding their care with less face-to-face contact.
- When in-person appointments are required (e.g. for blood tests, maternal examination or ultrasound scans), these should be arranged alongside other face-to-face maternity appointments to limit repeated clinic attendance.
- Particular consideration should be given to pregnant women who are 'shielding'. These women should be provided with a mask during hospital visits and shared waiting areas should be avoided.
  - If women who are 'shielding' attend hospital, they should be isolated in single rooms.
- Women should be able to notify the unit regarding self-isolation for COVID-19 using standard telephone numbers that are already available to them.
  - There should be a system in place to identify, support and follow up women who have missed appointments.

- o Units should appoint a group of clinicians to coordinate care for women forced to miss appointments due to self-isolation or a positive test. Missed appointments should be reviewed and either rescheduled if a face-to-face review is appropriate, converted to a remote appointment or deferred.
- For women receiving antenatal care across different sites, units must ensure that there are clear pathways for communication via handheld notes, electronic records and correspondence to general practitioners.
- Clinicians should be aware of specific changes to services which have been suggested via regularly updated, subspecialty service guidance available via the [RCOG website](#).

## Summary of evidence and rationale for guidance

Maternity care is essential, and studies in the UK and internationally have shown that women who do not attend antenatal services are at increased risk of maternal death, stillbirth and other adverse perinatal outcomes.<sup>37 38</sup> Antenatal and postnatal care should be therefore regarded as essential and women advised to attend, whilst observing current social distancing measures.

Recommendations on social distancing are available from the [UK Government](#).<sup>39</sup>

It is good practice to reconfigure services to reduce the number of times a woman is required to leave home to attend essential medical care. This may be achieved by offering consultations, investigations and care at the same clinic or hospital visit, or during virtual consultations or home visits where appropriate.<sup>40</sup> NHS England have issued [guidance](#) on the adoption of remote consultations in secondary care.<sup>41</sup>

The UK Government has published a list of conditions that make an [individual extremely vulnerable to the severe effects of COVID-19, along with guidance on how best to protect these individuals](#).<sup>42</sup>

Data on the effectiveness of telephone/video appointments are not available; until these are, we recommend clinicians follow locally agreed protocols for antenatal care delivery.

The care of pregnant women with complex healthcare needs is challenging during a pandemic. To support clinicians caring for these women, guidance documents to assist maternity units with changes to services were developed; these can be found on the [RCOG website](#) and are updated regularly.

## 2.2 What are the considerations for antenatal appointments?

### Advice

- Staff members should ensure adequate personal protective equipment (PPE) is used for face-to-face visits.
- Information and guidance should be available in languages spoken in the local communities served by the maternity unit.
- All women and any accompanying visitors (where permitted) should be advised to wear face masks or face coverings in line with national guidance.

- Evidence suggests that individuals of BAME background are at higher risk of developing severe complications of COVID-19. This appears to also apply to pregnant women. We therefore advise that:
  - Women of BAME background should be advised that they may be at higher risk of complications of COVID-19; we advise they seek advice without delay if they are concerned about their health.
  - Clinicians should be aware of this increased risk, and have a lower threshold to review, admit and consider multidisciplinary escalation of symptoms in women of BAME background.
  - When reorganising services, maternity units should be particularly cognisant of [evidence](#) that BAME individuals are at particular risk of developing severe and life-threatening COVID-19 disease.
  - Remote consultations should be encouraged, using interpreter services where appropriate, to minimise face-to-face contact.
- Women should be advised proactively to contact emergency antenatal services if they have any concern about their or their baby's wellbeing.
- Carbon monoxide testing of pregnant women has been paused during this period.
  - Midwives and doctors should still ask about and document smoking status at booking and 36 weeks, provide brief advice and refer women who smoke to specialist stop-smoking support on an opt-out basis.
- Women should continue to be encouraged to take folic acid and vitamin D supplements as per standard recommendations.
- Pregnant women will continue to need at least as much support, advice, care and guidance in relation to pregnancy, childbirth and early parenthood as before the pandemic, especially women living with adversity including poverty, homelessness, substance misuse, being an asylum seeker, experiencing domestic abuse and mental health problems.
  - Midwifery, obstetric and support staff should remain aware of the support needs for all women, acknowledging that trust-wide restrictions on hospital visitors are in place in most settings across the UK (including for women admitted to maternity services for antenatal and postnatal care) and may affect the amount of support that women require.
- Clinicians should be aware of the increased risk of domestic abuse in pregnancy, which has escalated during this pandemic. Women should be encouraged to share any concerns at every opportunity and provided with advice on how to access support if required.
- Women should be asked about their mental health at every contact. Women who require further support should be signposted to resources and local services, which may be remotely provided, where possible. These include:
  - [Sources of self-help](#) for anxiety and stress.

- When necessary, women in England can self-refer to local IAPT (Improving Access to Psychological Therapies) services.
- In Scotland, advice is available from [Parentclub](#) and [NHS Inform](#).
- Further information is available from the [Royal College of Midwives \(RCM\)](#) and [RCPsych websites](#).

## Summary of evidence and rationale for guidance

The appropriate use of PPE is an evolving area. We suggest that units follow the regularly updated [PHE guidance](#) on this in collaboration with their local guidance and infection control teams.<sup>43</sup> There are also clear guidelines on PPE from the [RCM](#).

The UK Government has issued guidelines on the use of face coverings within enclosed spaces in England, which apply to outpatient maternity appointments and hospital visitors.<sup>44</sup> We advise that these guidelines are followed. We will update if guidance relevant to the devolved nations becomes available.

Before this pandemic, there was already extensive evidence of the inequality of experience and outcomes for BAME women giving birth in the UK.<sup>25 26 45</sup> The increased risk of COVID-19 among BAME people is likely to result from a number of factors such as socioeconomic disadvantage, and the fact that they are more likely to work in key-worker roles including health and social care. Women of BAME background are disproportionately affected by particular inherited and metabolic conditions, and may avoid or delay seeking advice from health care professionals when unwell.<sup>46</sup> They should be encouraged to contact health services without delay. BAME women who are living with socioeconomic deprivation and/or in crowded conditions, those who were born outside the UK and whose first language is not English, and those with a high BMI and/or underlying medical conditions appear to be particularly high-risk groups.

There is currently an absence of accurate information about the additional risk of smoking and severe COVID-19 infection. This possible risk should be explained during smoking counselling in pregnancy.<sup>47</sup> The National Centre for Smoking Cessation and Training (NCSCT) has recommended that carbon monoxide testing during pregnancy be paused. Although manufacturers of carbon monoxide monitors support their safety in relation to viruses, in order to minimise face-to-face reviews and avoid workforce pressures, a pause in carbon monoxide monitoring is appropriate. Recommendations on smoking screening and cessation support are based on previous evidence on the effectiveness of these interventions. [Further guidance](#) is available, including in [Appendix H of the Saving Babies' Lives Care Bundle for England](#).<sup>48</sup>

This pandemic may result in an increased level of anxiety and other mental health problems in the general population.<sup>49</sup> There is increasing evidence that this is likely to be even greater for pregnant women as pregnancy represents a period of additional uncertainty.<sup>50 51</sup> Specifically, these anxieties are likely to revolve around: a) COVID-19 itself, b) the impact of social isolation resulting in reduced support from wider family and friends, c) the potential of reduced household finances and d) major changes in antenatal and other NHS care, including some appointments being changed from face-to-face to telephone contact.

Isolation, bereavement, financial difficulties, insecurity and inability to access support systems are all widely recognised risk factors for mental ill-health and are expected to affect individuals more than usual during the pandemic.<sup>50</sup>

[Recommendations on mental wellbeing](#) during the COVID-19 pandemic have been developed by the RCPsych and NHS England and Improvement.

The coronavirus pandemic has increased the incidence of domestic abuse.<sup>52,53</sup> Additional advice regarding support for victims of domestic abuse during the pandemic is [available here](#).

By acknowledging these difficulties, healthcare professionals can help to contain some of these anxieties.<sup>54</sup>

## **2.3 How should women with suspected or confirmed COVID-19 needing hospital attendance or advice be cared for?**

### **Advice**

For women who telephone maternity services:

- If women report symptoms attributed to COVID-19 on the phone to maternity services, consider differential diagnoses for fever, cough, change or loss of sense of smell/taste, or shortness of breath. This includes, but is not limited to urinary tract infection, chorioamnionitis and pulmonary embolism.
- If women have symptoms suggestive of COVID-19, they can self-refer to national services for [testing](#).

For women with possible or confirmed COVID-19, in whom hospital attendance is required or who self-present (this includes women who live with an individual who has possible or confirmed COVID-19):

- Women should be advised to attend via private transport where possible.
- If an ambulance is required, the call handler should be alerted if the woman, or a member of her household, is symptomatic of COVID-19.
- Women should be asked to alert a member of maternity staff by mobile telephone to their attendance when on the hospital premises, prior to entering the hospital or midwifery-led unit.
- Women should be met at the maternity unit entrance by staff wearing appropriate PPE, to provide a fluid-resistant surgical mask.
- Staff providing care should take PPE precautions as per national [health protection guidance](#).
- Women should immediately be escorted to an isolation room or cohort bay/ward, suitable for the majority of care during their hospital visit or stay.
  - Isolation rooms or ward bays should ideally have a defined area for staff to put on and remove PPE, and suitable bathroom facilities.
- The woman's face mask should not be removed until she is isolated in a suitable room or cohort bay.
- Only essential staff should enter the isolation room or bay.
- Visitors to isolation rooms or ward cohort bays should be kept to a minimum.

- All non-essential items from the isolation rooms should be removed prior to the woman's arrival (this includes other rooms in which the woman spends time during her hospital attendance [e.g. scan rooms if bedside scans are not appropriate]).
- All clinical areas must be cleaned following use, according to [current guidance](#).

## Summary of evidence and rationale for guidance

Availability of resources, provision of services and local prevalence of COVID-19 will vary across geographical regions, and will determine how women requiring hospital admission with confirmed or suspected COVID-19 are cared for. Advice on care in isolation rooms and COVID-19 cohort bays is [available from PHE](#).<sup>43</sup> This advice may change frequently and we urge clinicians to stay abreast of the latest developments.

As above, we suggest that units follow the regularly updated [PHE guidance](#) on PPE, in conjunction with guidance from the [RCM](#) and their local guidance and infection control teams.<sup>43 55</sup>

Guidance on cleaning clinical areas used to provide care to women with suspected or confirmed COVID-19 is available from PHE [health protection guidance](#).<sup>43</sup>

## 2.4 What are the considerations for antenatal care for women who have recovered from COVID-19?

### Advice

- For women who have recovered from COVID-19 with no, mild or moderate symptoms, without requiring admission to hospital, antenatal care should remain unchanged.
- Services should ensure that women who have missed antenatal appointments because of self-isolation are seen as early as is practical after the period of self-isolation ends.
- For women who have recovered from a period of serious or critical illness with COVID-19 requiring admission to hospital for supportive therapy, ongoing antenatal care should be planned together with a consultant obstetrician.
- Women who have been seriously or critically unwell should be offered a fetal growth scan approximately 14 days following recovery from their illness in the first instance, unless there is a pre-existing clinical reason for an earlier scan (e.g. fetal growth restriction [FGR]).

## Summary of evidence and rationale for guidance

Currently, there is an absence of evidence to guide the care for women recovering from mild or moderate symptoms of COVID-19. Women who have recovered should be encouraged to attend antenatal appointments in line with advice statements outlined in Section 2.4.

Although there is no evidence yet that FGR is a consequence of COVID-19, two-thirds of reported pregnancies with SARS were affected by FGR, so ultrasound follow-up seems prudent.<sup>56 57</sup>

Guidance on fetal growth surveillance following COVID-19 was developed along with the NHS England Saving Babies' Lives [modified Appendix G](#). This recommends a single fetal growth ultrasound scan a minimum of 14 days following resolution from acute illness of COVID-19 that required hospitalisation.





Royal College of  
Obstetricians &  
Gynaecologists

# 3. Venous thromboembolism prevention

## 3 Venous thromboembolism prevention

### 3.1 How should prevention of venous thromboembolism be addressed during the COVID-19 pandemic?

#### Advice

- Women who are self-isolating at home should stay well hydrated and mobile throughout this period.
- Women should have a venous thromboembolism (VTE) risk assessment performed during their pregnancy as per the RCOG Green-top Guideline No 37a. Infection with SARS-CoV-2 should be considered as a transient risk factor and prompt reassessment.
- Women already prescribed thromboprophylaxis should continue administering it.
- If women or healthcare professionals are concerned about the risk of VTE during a period of self-isolation, a clinical VTE risk assessment (in person or remotely) should be performed, and thromboprophylaxis considered and prescribed on a case-by-case basis.
- Local procedures should be followed in ensuring the supply of low molecular weight heparin (LMWH).
- Thromboprophylaxis commenced for pregnant women who are self-isolating should continue until they have recovered from the acute illness (between 7 and 14 days). For women with ongoing morbidity and limited mobility, advice from a clinician with expertise in VTE should be sought.
- All pregnant women admitted with confirmed or suspected COVID-19 should receive prophylactic LMWH, unless birth is expected within 12 hours.
- For women with severe complications of COVID-19, the appropriate dosing regimen of LMWH should be discussed in a multidisciplinary team (MDT) that includes a senior obstetrician or clinician with expertise in managing VTE in pregnancy.
- All pregnant women who have been hospitalised and have had confirmed COVID-19 should receive thromboprophylaxis for 10 days following hospital discharge. For women with persistent morbidity, consider a longer duration of thromboprophylaxis.
- If women are admitted with confirmed or suspected COVID-19 within 6 weeks postpartum, they should receive thromboprophylaxis for the duration of their admission and for at least 10 days post discharge. Consider extending this until 6 weeks postpartum for women with significant ongoing morbidity.

#### Summary of evidence and rationale for guidance

Pregnancy is widely recognised as a hypercoagulable state.<sup>58</sup> The existing RCOG Green-top Guidelines on VTE prevention and management should continue to support decision making during the COVID-19 pandemic.<sup>59</sup>

There is emerging evidence that suggests that individuals admitted to hospital with COVID-19 are also hypercoagulable.<sup>60</sup> Infection with SARS-CoV-2 is likely to be associated with an increased risk of maternal VTE. This is likely to be multifactorial, including the reduced mobility resulting from self-isolation at home or hospital admission, and other associated obstetric or maternal morbidity. Consequently, the cumulative risk is difficult to quantify.

The statements above were developed following expert consensus discussion to determine what increased risk COVID-19 may pose to pregnant women. VTE prevention for the unwell woman with COVID-19 is considered in section 5.2.



Royal College of  
Obstetricians &  
Gynaecologists

# 4. Labour and birth during the COVID-19 pandemic

## 4 Labour and birth during the COVID-19 pandemic

### 4.1 What are the considerations for labour and birth in asymptomatic women who test or have tested positive for SARS-CoV-2?

#### Advice

- For low-risk women who have no symptoms of COVID-19 but have tested positive for SARS-CoV-2 (within 7 days pre-birth) and wish to give birth at home or in a midwifery-led unit, it is recommended that an informed discussion around place of birth takes place with the midwife, consistent with local policies.
- For women who have no symptoms of COVID-19 but test positive for SARS-CoV-2 on admission, continuous electronic fetal monitoring (CEFM) during labour using cardiotocography (CTG) is not recommended routinely, unless it would normally be required for another reason (e.g. previous caesarean birth).
  - Fetal monitoring options should be discussed with the woman, acknowledging the evidence outlined below.
  - CEFM can be offered following a discussion about the current uncertainties in women who are asymptomatic with a positive test for SARS-CoV-2.

#### Summary of evidence and rationale for guidance

Whilst some case series have reported fetal compromise in women who are symptomatic of COVID-19, the need for CEFM for women who are asymptomatic of COVID-19 and otherwise low risk for labour (e.g. CEFM would not otherwise be indicated by NICE guidance on Intrapartum Care) is an area of clinical uncertainty due to the lack of robust evidence.<sup>61 62</sup> There is currently no evidence linking asymptomatic COVID-19 infection to abnormalities in continuous fetal monitoring or fetal compromise. While it is presently reassuring that there is no clear evidence of increased rates of fetal compromise in the asymptomatic population, women should continue to have the risks and benefits of CEFM discussed with them.

In the absence of other evidence, the current [NICE guidelines](#) should be followed.

### 4.2 How should a woman with suspected/confirmed COVID-19 be cared for in labour if they are symptomatic?

#### Advice

- Women with mild COVID-19 symptoms can be encouraged to remain at home (self-isolating) in early (latent phase) labour consistent with routine care. If women have symptoms suggestive of COVID-19, they can self-refer for [testing](#).

- If there are no concerns regarding the health of either the woman or baby, women who attend the maternity unit and would usually be advised to return home until labour is more established can still be advised to do so, unless private transport is not available.
  - Women should be given the usual advice regarding signs and symptoms of labour, but in addition should be informed about symptoms that might suggest deterioration related to COVID-19 and asked to call back if concerned.
- Advice on PPE is available in section 4.7.
- Women with symptomatic confirmed or suspected COVID-19 are recommended to labour and give birth in an obstetric unit.
- On admission, a full maternal and fetal assessment should be performed, including:
  - Assessment of the severity of COVID-19 symptoms by the most senior available clinician.
  - Maternal observations including temperature, respiratory rate and oxygen saturation.
  - Confirmation of the onset of labour, as per standard care.
  - CEFM using CTG.
- The following members of the MDT should be informed of the admission of the woman: consultant obstetrician, consultant anaesthetist, midwife-in-charge, consultant neonatologist and neonatal nurse in charge.
- Standard hourly maternal observations and assessment should be performed (as per the recommendations in the [NICE guideline on Intrapartum Care](#)), with the addition of hourly oxygen saturation.
- Oxygen should be titrated to aim for saturation above 94%.
- Women with symptomatic confirmed or suspected COVID-19 should be offered CEFM during labour and vaginal birth.
- Efforts should be made to minimise the number of staff members entering the room and units should develop a local policy specifying essential personnel for emergency scenarios.
- Water births are not recommended due to the potential risk of disease transmission through faeces.

## Summary of evidence and rationale for guidance

NHS England has also produced clinical guidance on the [temporary reorganisation of intrapartum maternity care during the coronavirus pandemic](#).<sup>64</sup>

PHE COVID-19 [infection and control guidance](#) gives advice about avoiding disease transmission.<sup>43</sup>

[WHO](#) have produced guidance on clinical management of COVID-19.

In women with symptomatic COVID-19, there may be an increased risk of fetal compromise in active labour.<sup>61 62</sup> Although the data in this area are poor, it appears prudent to use fetal monitoring for maternal systemic infection including COVID-19.

Many of the advice statements above are recommendations of good practice points derived from expert consensus.

Women have been advised to avoid water births due to the risk of disease transmission through faeces, although studies are in small numbers with possibly low viral loads in stool samples.<sup>65 66</sup>

### **4.3 What are the considerations for labour and birth for women who have recovered from COVID-19?**

#### **Advice**

- For women who have recovered from COVID-19, without requiring admission to hospital, and who have completed self-isolation in line with public health guidance, there should be no change to planned care during labour and birth.
- For women who have recovered following a hospital admission for serious or critical COVID-19 illness needing supportive therapy, place of birth should be discussed and planned with the woman, her family, if she wishes, and a consultant obstetrician. A personalised assessment should take into consideration fetal growth and the woman's choices.
- Informed discussions with women about fetal monitoring should acknowledge that evidence of fetal distress is based on small numbers of babies and theoretical risks extrapolated from pregnancies affected by FGR in women with other coronaviruses.

#### **Summary of evidence and rationale for guidance**

There is an absence of evidence for this situation. The above is based on expert consensus.

### **4.4 What about birth partners during the COVID-19 pandemic?**

#### **Advice**

- Women should be supported and encouraged to have birth partners present with them during active labour and birth if they wish to do so, unless the birth occurs under general anaesthetic (GA), in accordance with local or national hospital policies.
- If symptomatic or in a period of [self-isolation](#) for confirmed SARS-CoV-2 infection, birth partners should remain in self-isolation at home and not attend the unit.
- On attendance at the maternity unit, all birth partners should also be asked whether they have had any symptoms suggestive of COVID-19 – e.g. fever, acute persistent cough, hoarseness, anosmia, nasal discharge/congestion, shortness of breath, sore throat, changes in or loss of sense of smell or taste, wheezing or sneezing, in the preceding 7 days.

- If these symptoms began within 7 days or less, or they remain symptomatic (other than with a persistent cough), they should be asked to leave the maternity unit immediately and self-isolate at home.
- Guidance about testing for women and their birth partners is discussed in the [RCOG document](#) on women seeking maternity care in a hospital setting.
- We recommend that asymptomatic birth partners are permitted to stay with the woman through labour and birth, unless the birth occurs under general anaesthetic.
- Birth partners should be asked to remain by the woman's bedside and to not walk around the ward/hospital.
- Restrictions on other visitors should follow local hospital policy.

### Summary of evidence and rationale for guidance

Having a trusted birth partner present throughout labour is known to make a significant difference to the safety and wellbeing of women in childbirth.<sup>67-69</sup>

General PHE guidance, local hospital infection control and visitor policies should be adhered to.<sup>43 70</sup>

## 4.5 What informed discussions should take place with women regarding timing and mode of birth during the COVID-19 pandemic?

### Advice

- Information regarding mode of birth during the COVID-19 pandemic should be discussed with the woman and her family, taking into consideration her preferences and any obstetric indications for intervention.
- Guidance on testing all women attending maternity services is summarised in the [RCOG's 'Principles for the testing and triage of women seeking maternity care in hospital settings, during the COVID-19 pandemic'](#).
- A personal assessment should be made to determine whether it is beneficial overall to delay elective caesarean birth or induction of labour (IOL), and any associated appointments, for women who are currently in a period of self-isolation because of suspected COVID-19 in themselves or a household contact.
  - Personalised assessments to consider delaying elective birth for women in self-isolation should take into account the urgency of the birth and the risk of infectious transmission to other women, healthcare workers and, postnatally, to her baby.
  - If a planned caesarean birth or IOL cannot be delayed, the advice for services providing care to women admitted when affected with suspected/confirmed COVID-19 should be followed.
- The use of birthing pools in hospital should be avoided in suspected or confirmed cases of COVID-19, given the potential risk of infection via faeces.



- In women with symptoms who are becoming exhausted or hypoxic, a personalised and informed discussion and decision should be made regarding shortening the length of the second stage of labour with instrumental birth.
- When urgent birth of the baby is required to aid supportive care of a woman with severe or critical COVID-19 and vaginal birth is not imminent, consider whether the benefits of an urgent caesarean birth may outweigh any risks to the woman.
- The advice in section 4.7 should be followed for PPE for caesarean birth.
- Women and their families should be aware that donning PPE for emergency caesarean births is time-consuming but essential, and that this may impact on the time to delivery and potentially result in adverse outcome. This should be taken into account during decision making.

### Summary of evidence and rationale for guidance

There is no evidence to favour one mode of birth over another in women with COVID-19. There is currently limited direct evidence to demonstrate intrauterine vertical transmission of SARS-CoV-2.<sup>71</sup> In the UKOSS study, 12 (5%) babies tested positive for SARS-CoV-2 infection; six within the first 12 hours (two were born by unassisted vaginal birth and four by caesarean birth) and six after 12 hours (two born vaginally and four by caesarean birth).<sup>13</sup>

## 4.6 What are the specific considerations for labour analgesia or anaesthesia?

### Advice

- Entonox can be used with a single-patient microbiological filter.
- In early labour, inform women with symptomatic or confirmed COVID-19 about the potential benefit of epidural analgesia in minimising the need for GA if urgent intervention for birth is needed, so that they can make informed decisions regarding use or type of labour analgesia/anaesthesia.

### Summary of evidence and rationale for guidance

Advice published on the considerations for [labour analgesia or anaesthesia](#) is based on expert opinion following consultation with the Obstetric Anaesthetists' Association.

There is no evidence that the use of Entonox is an [aerosol-generating procedure](#) (AGP).

There is no evidence that [epidural or spinal analgesia or anaesthesia](#) is contraindicated in the presence of coronaviruses.<sup>72</sup>

Intubation, required for GA in the case of caesarean birth, is an [AGP](#). This significantly increases the risk of transmission of coronavirus to the attending staff.

## 4.7 What personal protective equipment is recommended when caring for women during labour and birth?

### Advice

- Guidance on PPE in all scenarios, including during labour and birth, has been issued by PHE.<sup>43</sup>
- The level of PPE required for caesarean birth requires a multidisciplinary discussion about the likelihood of requiring a GA.
- For the minority of caesarean births, where GA is planned from the outset, all staff in theatre should wear PPE, including an FFP3 mask and visor. PPE should be donned prior to commencing the GA.
- Local standard operative procedures should be developed to determine type of PPE for cases of unsuccessful regional anaesthesia.

### Summary of evidence and rationale for guidance

General advice from PHE on type and specification of PPE is [available here](#). Particular advice from PHE on type and specification of PPE for different maternity settings is available as part of the [table here](#).

The level of PPE required by healthcare professionals caring for a woman with COVID-19 who is undergoing a caesarean birth should be determined on the basis of the risk of requiring a GA. Intubation is an [AGP](#). This significantly increases the risk of transmission of coronavirus to the attending staff.

Siting regional anaesthesia (spinal, epidural or combined spinal epidural [CSE]) is not an AGP.

The chance of requiring conversion to a GA during a caesarean birth commenced under regional anaesthesia is small, but this chance increases with the urgency of caesarean birth. In situations where there are risk factors that make conversion to a GA more likely, the decision on what type of PPE to wear should be judged on basis of the individual circumstances. If the risk of requiring conversion to a GA is considered significant (e.g. 'top-up' of a suboptimal epidural from labour), the theatre team should wear PPE appropriate to a GA in readiness.

## 4.8 How should obstetric theatres be managed during the COVID-19 pandemic?

### Advice

- Elective obstetric procedures (e.g. cervical cerclage or caesarean birth) planned for women with suspected/confirmed COVID-19 should be scheduled at the end of the operating list.
- Emergency procedures for women with suspected/confirmed COVID-19 should be carried out in a second obstetric theatre, where available, allowing time for a full postoperative theatre clean as per national health protection guidance.
- The number of staff in the operating theatre should be kept to a minimum, and all must wear appropriate PPE.

- Anaesthetic care for women with suspected or confirmed COVID-19 should be with reference to guidance from Royal College of Anaesthetists/Obstetric Anaesthetists' Association.
- The use of PPE causes communication difficulties in obstetric theatres so checklists should be used with closed loop communication.<sup>73</sup>

### Summary of evidence and rationale for guidance

The statements above were developed on the basis of government advice [on IPC](#).<sup>43</sup>

The guidance above is based on expert consensus and IPC advice.



Royal College of  
Obstetricians &  
Gynaecologists

# 5. Managing clinical deterioration during the COVID-19 pandemic

## 5. Managing clinical deterioration during the COVID-19 pandemic

### 5.1 How should a woman requiring hospital admission with symptoms suggestive of COVID-19 be investigated?

#### Advice

- If the woman attends with a fever, investigate and treat as per [RCOG guidance on sepsis in pregnancy](#). Testing for COVID-19 should be arranged in addition to blood cultures.
- While pyrexia may suggest COVID-19, do not assume that all pyrexia is due to COVID-19. Consider the possibility of bacterial infection and perform full sepsis-six screening and administer intravenous antibiotics when appropriate.
- Consider bacterial infection if the white blood cell count is raised (lymphocytes usually normal or low with COVID-19) and commence antibiotics.
- Women should be tested for COVID-19 if they meet the inpatient or community [PHE criteria](#).
  - Current inpatient case criteria (correct as of 2 June 2020) are individuals who are being/are admitted to hospital with one of the following:
    - A loss of, or change in, normal sense of taste or smell (anosmia) in isolation or in combination with any other symptoms of COVID-19.
    - Clinical/radiological evidence of pneumonia.
    - ARDS.
    - Fever  $\geq 37.8^{\circ}\text{C}$  AND at least one of the following: acute persistent cough, hoarseness, nasal discharge/congestion, shortness of breath, sore throat, wheezing or sneezing.
- Radiographic investigations should be performed as for the non-pregnant adult; this includes chest X-ray and computerised tomography (CT) of the chest.
  - Chest imaging, especially CT of the chest, is essential for the evaluation of the unwell woman with COVID-19 and should be performed when indicated, and not delayed because of concerns of possible fetal exposure to radiation, as maternal wellbeing is paramount.
- The diagnoses of pulmonary embolism and heart failure should be considered in women with chest pain, worsening hypoxia or a respiratory rate  $>22$  breaths/min (particularly if there is a sudden increase in oxygen requirements), or in women whose breathlessness persists or worsens after expected recovery from COVID-19.

- Consider additional investigations to rule out differential diagnoses – e.g. electrocardiogram (ECG), CT pulmonary angiogram, echocardiogram, etc.

## Summary of evidence and rationale for guidance

The clinical symptoms of COVID-19 overlap with those of a variety of other clinical conditions. We encourage clinicians to consider all differential diagnoses for women who present with a fever in pregnancy and follow the advice and guidance of the RCOG Green-top Guideline No. 64a.<sup>74</sup>

The current PHE guidance on testing for suspected COVID-19 is [available here](#).<sup>16</sup> This is a rapidly evolving area of policy and will likely have local variations on implementation. We recommend that clinicians liaise with their local infection control and testing teams. In addition, a [national framework](#) that applies to NHS Trusts in England at present recommends offering testing for SARS-CoV-2 to all hospital inpatients. [Specific guidance from the RCOG](#) has been developed that recommends how this framework may need to be modified for maternity services.

Several studies have shown decreased lymphocyte counts in the general population affected by COVID-19.<sup>75</sup> One systematic review noted decreased lymphocyte counts in 164 pregnant women.<sup>76</sup>

## 5.2 How should a woman with suspected/confirmed COVID-19 who is clinically deteriorating be cared for?

### Advice

- Obstetricians should be familiar with local protocols for the initial investigation and care of patients presenting to medical teams with possible COVID-19. These protocols should be followed for pregnant women as far as possible (including initial investigations, management of fluid balance and escalation of care to involve the critical care team).
- The priority for medical care should be to stabilise the woman's condition with standard therapies.
- Hourly observations should include respiratory rate and oxygen saturation, monitoring both the absolute values and trends.
- Signs of decompensation include an increase in oxygen requirements or  $FiO_2 > 40\%$ , a respiratory rate  $>30/\text{min}$ , reduction in urine output, or drowsiness, even if oxygen saturations are normal.
- Escalate urgently if any signs of decompensation develop.
  - Young, fit women can compensate for a deterioration in respiratory function and are able to maintain normal oxygen saturations until sudden decompensation.
  - Signs of decompensation include an increase in oxygen requirements, an increasing respiratory rate despite oxygen therapy, an acute kidney injury or drowsiness even if the saturations are normal.
- Titrate oxygen flow to maintain saturations  $>94\%$ .

- Have a low threshold to start antibiotics at presentation, with early review and rationalisation of antibiotics if COVID-19 is confirmed. Even when COVID-19 is confirmed, remain open to the possibility of another co-existing condition.
- Suspected COVID-19 should not delay administration of therapy that would usually be given (e.g. intravenous antibiotics in woman with fever and prolonged rupture of membranes).
- Until test results are available, a woman with suspected COVID-19 should be treated as though it is confirmed.
- An MDT planning meeting should be urgently arranged for any unwell woman with suspected/confirmed COVID-19. This should ideally involve a consultant physician, consultant obstetrician, midwife-in-charge, consultant neonatologist, consultant anaesthetist and intensivist responsible for obstetric care. The discussion should be shared with the woman and her family if she chooses. The following considerations should be included:
  - Key priorities for medical care of the woman and her baby, and her birth preferences.
  - Most appropriate location of care (e.g. intensive care unit, isolation room in infectious disease ward or other suitable isolation room) and lead specialty.
  - Concerns among the team regarding special considerations in pregnancy, particularly the health of the baby.
- All pregnant women should have a VTE assessment and be prescribed prophylactic dose thromboprophylaxis, unless there is a suspicion of a VTE when therapeutic dose thromboprophylaxis should be administered.
- For women with thrombocytopenia (platelets <50), stop aspirin prophylaxis and thromboprophylaxis and seek haematology advice.
- Be aware of possible myocardial injury, and that the symptoms are similar to those of respiratory complications of COVID-19.
- Be aware of the interim government guidance based on the results of the RECOVERY trial, which states that steroid therapy should be considered for 10 days or to hospital discharge, whichever is sooner, for adults unwell with COVID-19 and requiring oxygen (in pregnant adults, use oral prednisolone 40 mg once a day or intravenous hydrocortisone 80 mg twice a day).
- Apply caution with intravenous fluid management:
  - Women with moderate-to-severe symptoms of COVID-19 should be monitored using hourly fluid input/output charts.
  - Efforts should be targeted towards achieving neutral fluid balance in labour.
  - Try boluses in volumes of 250–500 ml and then assess for fluid overload before proceeding with further fluid resuscitation.

- The frequency and suitability of fetal heart rate monitoring should be considered on an individual basis, accounting for the gestational age and the maternal condition.
- An individualised assessment of the woman should be made by the MDT to decide whether emergency caesarean birth or IOL is indicated, either to assist efforts in maternal resuscitation or where there are serious concerns regarding the fetal condition.
  - Individual assessment should consider: the maternal condition (including changes in oxygen saturations, radiological changes and respiratory rate), the fetal condition, the potential for improvement following iatrogenic birth, and the gestation. The priority must always be the wellbeing of the woman.
  - If urgent intervention for birth is indicated for fetal reasons, birth should be expedited as for normal obstetric indications, as long as the maternal condition is stable.
  - If maternal stabilisation is required before intervention for birth, this is the priority, as it is in other maternity emergencies (e.g. severe pre-eclampsia).
- Antenatal steroids for fetal lung maturation should be given when indicated by NICE guidance but urgent intervention for birth should not be delayed for their administration.
- Consider administering magnesium sulphate cover for fetal neuroprotection irrespective of steroid status, but do not delay to administer the magnesium sulphate if urgent birth is indicated.

## Summary of evidence and rationale for guidance

A useful summary on supportive care for adults diagnosed with COVID-19 has been published by [WHO](#).<sup>77</sup> Specific guidance on the management of patients with COVID-19 who are admitted to critical care has now been published by [NICE](#) and [SIGN](#).<sup>78,79</sup>

In a retrospective observational study of 12 pregnant women with severe disease from COVID-19, nine needed respiratory support; eight of these women were delivered with maternal respiratory distress. Seven of those did not require intubation, with two of them having improved oxygenation within 2 hours postpartum, showing that birth did not worsen the respiratory status.<sup>80</sup>

Severe COVID-19 may be associated with thrombocytopenia. When aspirin has been prescribed as prophylaxis for pre-eclampsia, it should be discontinued as this may increase the bleeding risk in thrombocytopenic patients.<sup>81</sup>

Myocardial injury and its complications were observed in 9.5% of all patients who died in Italy up to 13 April 2020.<sup>82</sup> Early involvement of multidisciplinary colleagues to investigate for potential myocardial injury is essential if this is suspected.<sup>83</sup> Further details of investigation and management is available in the [NICE rapid guideline](#) on diagnosing myocardial injury in patients with suspected or confirmed COVID-19.<sup>83</sup>

Antenatal corticosteroids are well established as being beneficial in threatened preterm labour, or if iatrogenic preterm birth is anticipated.<sup>84</sup> There is no evidence that steroids in the doses prescribed for fetal lung maturation cause any harm in the context of COVID-19.<sup>85</sup> Magnesium sulphate is



recommended for fetal neuroprotection in preterm babies as per RCOG guidelines.<sup>86</sup>

For non-specialist anaesthetists and physicians involved in the care of pregnant women with COVID-19 and other medical conditions, useful information is available from the guidelines on '[Care of the Critically Ill Woman in Childbirth](#)' and the Royal College of Physicians' [Acute Care toolkit 15: managing acute problems in pregnancy](#).<sup>87 88</sup>

The interim results of the RECOVERY trial demonstrated a significant reduction in mortality for individuals with COVID-19 requiring oxygen who were given steroid therapy.<sup>89</sup> This has immediately been recommended for use in the NHS.<sup>90</sup> The RECOVERY trial protocol for pregnancy recommends prednisolone 40 mg orally once daily, and in women unable to take oral medicine, hydrocortisone 80 mg intravenously twice daily instead of dexamethasone treatment.<sup>91</sup>



Royal College of  
Obstetricians &  
Gynaecologists

## 6. Postnatal care

## 6. Postnatal care

Routine postnatal care for women in accordance with [national guidelines](#) and the [RCOG framework for service modifications to antenatal and postnatal care](#) during the pandemic should be followed.

### 6.1 How should neonatal care for the baby be provided during the COVID-19 pandemic?

#### Advice

- Women and their healthy babies, who do not otherwise require maternal critical care or neonatal care, should be kept together in the immediate postpartum period.
- A risks and benefits discussion with neonatologists and families is recommended to individualise care in babies who may be more susceptible.
- All babies born to SARS-CoV-2-positive women should be cared for as per [guidance from the Royal College of Paediatrics and Child Health \(RCPCH\)](#).<sup>92</sup>
- Specific guidance on neonatal resuscitation during the COVID-19 pandemic is available from the [Resuscitation Council](#).

#### Summary of evidence and rationale for guidance

There are limited data to guide the neonatal care of babies of women who tested positive for SARS-CoV-2 in the third trimester.<sup>93 94</sup>

### 6.2 What should parents/carers be advised regarding infant feeding during the COVID-19 pandemic?

#### Advice

- Parents should be supported in their feeding choices and informed of the risks and benefits of feeding the baby in close proximity, particularly to individuals with suspected or confirmed COVID-19.
- Parents should be informed that infection with COVID-19 is not a contraindication to breastfeeding.
- The following precautions should be taken to limit viral spread to the baby:
  - Considering asking someone who is well to feed the baby.
  - Wash hands before touching the baby, breast pump or bottles.

- Avoid coughing or sneezing on the baby while feeding.
- Consider wearing a face covering or fluid-resistant face mask while feeding or caring for the baby.
- Babies should not wear masks or other face coverings, as they may risk suffocation.
- When women are expressing breastmilk in hospital, a dedicated breast pump should be used.
  - Where a breast pump is used, follow recommendations for pump cleaning after each use.
  - For babies who are bottle-fed with formula or expressed milk, strict adherence to [sterilisation guidelines](#) is recommended.

## Summary of evidence and rationale for guidance

The benefits of breastfeeding outweigh any potential risks of transmission of the virus through breastmilk. It is reassuring that a recent systematic review found that, in 24 cases, breastmilk tested negative for COVID-19; however, given the small number of cases, this evidence should be interpreted with caution.<sup>14</sup> The main risk of breastfeeding is the close contact between the baby and the woman, who is likely to share infective droplets.

In light of the current evidence, we advise that the benefits of breastfeeding outweigh any potential risks of transmission of the virus through breastmilk.<sup>95</sup> This is a view supported by the [UNICEF Baby Friendly Initiative](#), which is widely implemented in the UK.<sup>96</sup>

Specific recommendations on minimising the risk of transmission when feeding the baby were developed with experts from RCPCH, and from their guidance.<sup>92</sup>

Face coverings are not deemed appropriate for babies. The current government advice for using face coverings is directed towards adults in England.<sup>97</sup>

### 6.3 What are the considerations for postnatal care for women and babies following admission with COVID-19?

#### Advice

- All households are advised to self-isolate at home for 14 days after birth of a baby to a woman with current COVID-19.
- Usual advice about safe sleeping and a smoke-free environment should be emphasised, along with provision of clear advice about careful hand hygiene and infection control measures when caring for and feeding the baby.
- Families should be provided with guidance about how to identify signs of illness in their newborn or worsening of the woman's symptoms, and provided with appropriate contact details if they have concerns or questions about their baby's wellbeing.

- Any women or babies requiring readmission for postnatal obstetric or neonatal care during a period of self-isolation for suspected or confirmed COVID-19 are advised to telephone their local unit ahead of arrival.

### Summary of evidence and rationale for guidance

The RCPCH has published guidance on the neonatal care of babies born to women with COVID-19.<sup>92</sup> The advice for households to isolate for 14 days after the birth of a baby born to a woman who is infected with SARS-CoV-2 is to ensure a full period of isolation in case of incubation of the virus in the baby. These advice statements have been extrapolated from the RCPCH guidance and developed using expert consensus opinion.

# Acknowledgments

The RCOG COVID-19 Guidance Cell is comprised of: Dr Eddie Morris (President, RCOG), Professor Tim Draycott (Vice President for Clinical Quality, RCOG), Dr Pat O'Brien (Vice President for Membership, RCOG), Anita Powell (Senior Director for Clinical Quality, RCOG), Dr Mary Ross-Davie (Director for Scotland, RCM), Dr Jennifer Jardine (Clinical Fellow, RCOG), Dr Sophie Relph (Clinical Fellow, RCOG), Dr Gemma Goodyear (Clinical Fellow, RCOG), Dr Jahnavi Daru (Honorary Clinical Fellow, RCOG), Dr Christine Ekechi (Honorary Clinical Fellow, RCOG), Dr Anushka Tirlapur (Honorary Clinical Fellow, RCOG), Gemma Thurston (Business Manager, RCOG), Louise Thomas (Head of Quality Improvement, RCOG), Emma Gilgunn-Jones (Director of Media and PR, RCOG), Jenny Priest (Director of Policy and Public Relations, RCOG), Gozde Zorlu (Media and PR Manager, RCOG) and Stephen Hall (Political Advisor, RCOG).

The following external subject experts contributed to the sections on VTE (Professor Beverley Hunt, Professor Catherine Nelson-Piercy, Professor Rezan Abdulkadir, Dr Peter MacCallum, Dr Louise Bowles and Dr Shohreh Beski) and Managing Clinical Deterioration of COVID-19 (Professor Catherine Nelson-Piercy, Dr Margaret Blott, Dr Arlene Wise and Professor Lucy Chappell).

We also wish to acknowledge the contributions of Professor Russell Viner and Dr David Evans (on behalf of the RCPCH), Dr Fiona Donald and Dr Nuala Lucas (on behalf of the Obstetric Anaesthetists' Association and Royal College of Anaesthetists'), Dr Giles Berrisford (on behalf of RCPsych), colleagues from the digital and communications teams at RCOG and the following individual contributors: Professor Asma Khalil, Dr Lucy Mackillop, Dr Charlotte Frise, Dr Toni Hazell, Dr Ed Mullins, Dr Benjamin Black and Zeenath Uddin.

Finally, we wish to acknowledge the rapid peer-review of the following individuals and organisations: Dr Matthew Jolly and colleagues at NHS England, Dr Corinne Love and colleagues at the Scottish Government, Emma Crookes (RCOG Women's Voices), the British Intrapartum Care Society (BICS), the British Association of Perinatal Medicine (BAPM), the British Maternal and Fetal Medicine Society (BMFMS) and members of the RCOG Guidelines Committee.



Royal College of  
Obstetricians &  
Gynaecologists

# Appendix

## Appendix I: Summary of previous updates

Version	Date	Summary of changes
<b>2</b>	12.3.20	<b>1.2:</b> At the time of writing, Public Health Wales are aligning with Public Health England on case definitions, assessment, infection prevention and control and testing. We will update <a href="#">this guidance</a> if this changes.
<b>2</b>	13.3.20	<b>2.2:</b> Updated to reflect PHE and health protection advice as per 13.03.20, in particular to use online symptom checkers and to treat all individuals with symptoms as possibly having COVID-19.
<b>2</b>	13.3.20	<b>3.2:</b> Sentence on who to test updated to reflect advice to test women with symptoms suggestive of COVID-19 who require admission.
<b>2</b>	13.3.20	<b>3.6.4 and 3.6.5:</b> Updated to suggest considering delay of elective caesarean birth or induction for women with symptoms suggestive of COVID-19 as well as those with confirmed COVID-19.
<b>2</b>	13.3.20	<b>3.8:</b> Infant feeding modified from recommendation to wear a face mask to try and avoid coughing or sneezing on the baby, and consider wearing face mask where available.
<b>2</b>	13.3.20	<b>4:</b> New section added for antenatal care for pregnant women following self-isolation for symptoms suggestive of COVID-19.
<b>2</b>	13.3.20	<b>5 (new).</b> New section - Advice for pregnant healthcare professionals.
<b>2</b>	13.3.20	<b>Appendix I:</b> Flow chart amended to reflect modified PHE guidance.
<b>2</b>	13.3.20	<b>References: 19:</b> NHS Staff Council Statement on Covid-19 2020 [Available from: <a href="https://www.nhsemployers.org/-/media/Employers/Documents/Pay-and-reward/NHS-Staff-Council---Guidance-for-Covid-19-Feb-20.pdf?la=en&amp;hash=70C909DA995280B9FAE4BF6AF291F4340890445C">https://www.nhsemployers.org/-/media/Employers/Documents/Pay-and-reward/NHS-Staff-Council---Guidance-for-Covid-19-Feb-20.pdf?la=en&amp;hash=70C909DA995280B9FAE4BF6AF291F4340890445C</a> ] accessed 12 March 2020.
<b>3</b>	17.3.20	<b>2:</b> Advice for Health Professionals to share with Pregnant Women updated to reflect current guidelines.
<b>3</b>	17.3.20	<b>3:</b> New section added on Advice for all midwifery and obstetric services.
<b>3</b>	17.3.20	<b>4.1:</b> General advice to services providing care to pregnant women updated to reflect advice from chief medical officer on 16/3/20.
<b>3</b>	14.3.20	<b>4.1:</b> Advice on cleaning ultrasound equipment added, and reference added.
<b>3</b>	17.3.20	<b>4.5:</b> Linked to new national guidance on the actions required when a COVID-19 case was not diagnosed on admission .
<b>3</b>	17.3.20	<b>4.6.2:</b> Recommendations added: There is evidence of household clustering and household co-infection. Asymptomatic birth partners should be treated as possibly infected and asked to wear a mask and wash their hands frequently. If symptomatic, birth partners should remain in isolation and not attend the unit.  The use of birthing pools in hospital should be avoided in suspected or confirmed cases, given evidence of transmission in faeces and the inability to use adequate protection equipment for healthcare staff during water birth.



<b>3</b>	17.3.20	<b>4.6.2:</b> Advice about Entonox changed to  There is no evidence that the use of Entonox is an aerosol-prone procedure  Entonox should be used with a single-patient microbiological filter. This is standard issue throughout maternity units in the UK.
<b>3</b>	17.3.20	<b>4.6.4:</b> Anaesthetic management for women with symptoms or confirmed COVID-19, which was previously in this guidance, has been removed and external links provided
<b>3</b>	17.3.20	<b>4.7.1:</b> Statement inserted 'Chest imaging, especially CT chest, is essential for the evaluation of the unwell patient with COVID-19 and should be performed when indicated and not delayed due to fetal concerns.'
<b>3</b>	17.3.20	Updated to reflect current public health guidance on self-isolation and social distancing.
<b>3</b>	17.3.20	<b>4.7.1:</b> Advice on neonatal management and testing has been removed. Please refer to <a href="#">RCPCH guidance</a> .
<b>3</b>	17.3.20	<b>6:</b> Advice for healthcare professionals updated in line with Chief Medical Officer statement on Monday 16 March.
<b>4</b>	21.3.20	<b>6:</b> Section on 'Occupational health advice for employers and pregnant women during the COVID-19 pandemic' added, replacing the previous section 6 on 'Information for Healthcare Professionals'. Section includes specific recommendations for healthcare professionals.
<b>4</b>	21.3.20	<b>1.3-1.4:</b> Additional information added on the susceptibility of pregnant women to COVID-19 infection.
<b>4</b>	21.3.20	<b>2:</b> Additional information on social distancing for pregnant women added, particularly specifying stringent adherence to recommendations for women >28 weeks gestation.
<b>4</b>	21.3.20	<b>4.7:</b> New section added on specific recommendations for PPE during labour and birth.
<b>4</b>	21.3.20	<b>1:</b> Addition of information and links for the UKOSS reporting system.
<b>4</b>	21.3.20	<b>All:</b> General proofread and editorial changes.
<b>4</b>	21.3.20	<b>6:</b> Page 36 title changed to 'Occupational health advice for employers and pregnant women during the COVID-19 pandemic'.
<b>4.1</b>	26.3.20	<b>Chapter 6:</b> 'Occupational health advice for employees and pregnant women during the COVID-19 pandemic' has been removed from this general guidance on pregnancy and COVID-19 infection, and published as a separate document given the distinct audience for the occupational health advice.
<b>4.1</b>	26.3.20	<b>4.7.3:</b> On Personal Protective Equipment updated in line with NHS England guidance.

<b>5</b>	28.3.20	<b>1.3:</b> Section updated to include new evidence on possible vertical transmission.
<b>5</b>	28.3.20	<b>2.2:</b> Sentence added on the major new measures announced by government for pregnant women with co-existing significant congenital or acquired heart disease.
<b>5</b>	28.3.20	<b>2.3:</b> Section updated to emphasise the need to attend maternity care.
<b>5</b>	28.3.20	<b>3:</b> General advice for antenatal care extended to include considerations for vulnerable women. Section also added on general advice regarding intrapartum services.
<b>5</b>	28.3.20	<b>3.1:</b> Specific advice added regarding the cessation of carbon monoxide monitoring in pregnancy, following advice from the National Centre for Smoking Cessation and Training.
<b>5</b>	28.3.20	<b>4:</b> Scotland specific links to Health Protection Scotland removed after confirmation from the Scottish government that National links from gov.uk should be used.
<b>5</b>	28.3.20	<b>4.3.6:</b> Scotland specific links to Health Protection Scotland removed after confirmation from the Scottish government that National links from gov.uk should be used.
<b>5</b>	28.3.20	<b>4.7.3 and 4.76:</b> Advice on PPE considerations for caesarean birth and general advice for obstetric theatres moved to new section 'Specific peri-operative advice for pregnant women with suspected/confirmed COVID-19 requiring surgical intervention'.
<b>5</b>	28.3.20	<b>4.8.1:</b> Reference made to new guidance published by NICE on the management of patients with COVID-19 in critical care.
<b>5</b>	28.3.20	<b>4.8.1:</b> Additional recommendations made for the management of women admitted during pregnancy with suspected/confirmed COVID-19.
<b>5</b>	28.3.20	<b>4.9.2:</b> Section edited to make infant feeding recommendations to any caregiver; not just to the mother.
<b>5</b>	28.3.20	<b>4.10:</b> New section on 'Specific peri-operative advice for pregnant women with suspected/confirmed COVID-19 requiring surgical intervention'.
<b>5</b>	28.3.20	<b>5.1:</b> Correction of an error in the title to clarify that this section refers to the care of women recovering from suspected (not confirmed) COVID-19 for which hospitalisation was not required.
<b>6</b>	3.4.20	<b>Throughout:</b> References to the new RCOG guidance on (1) antenatal and postnatal services (2) antenatal screening (3) fetal medicine services (4) maternal medicine services and (5) self-monitoring of blood pressure, have been added throughout the document.
<b>6</b>	3.4.20	<b>1.2:</b> New resources signposted on current UK and international disease incidence.
<b>6</b>	3.4.20	<b>1.4:</b> Sentence reporting that there are 'no reported maternal deaths from COVID-19' removed because there was recently a possible maternal death reported in tabloid media. There is not any robust evidence to amend this statement or report confidently in the guideline.
<b>6</b>	3.4.20	<b>3.2:</b> Addition of new advice on screening birth partners for recent possible symptoms of COVID-19 when they attend the maternity unit. In addition, suggestion of information to give the birth partner about what is expected of them whilst they are in the hospital, to assist staff in reducing the risk of infection transmission and to assist with communication when birth partners accompany women into operating theatres.
<b>6</b>	3.4.20	<b>3.4:</b> Moved to section 3.2.
<b>6</b>	3.4.20	<b>3.5:</b> New section on maternal mental wellbeing during the pandemic.
<b>6</b>	3.4.20	<b>4.1</b> The previous section 4.2 was repetitive of section 3.1 and so has been removed. Sections 4.2 onwards have been re-numbered.

<b>6</b>	3.4.20	<b>4.3:</b> Inclusion of the PHE case definition for COVID-19 testing, rather than referring readers to this through the link.
<b>6</b>	3.4.20	<b>4.9:</b> Updates to advice on PPE for caesarean birth, to ensure that these are consistent with new PHE advice.
<b>7</b>	9.4.20	<b>1.4:</b> Update to data from ICNARC and inclusion of a report of 43 pregnant women with COVID-19 from New York.
<b>7</b>	9.4.20	<b>1.4:</b> New comment on risk of venous thromboembolism from COVID-19.
<b>7</b>	9.4.20	<b>2.3:</b> Advice for pregnant women added – if they are advised to attend a face-to-face antenatal appointment, this is because the appointment is important and the benefit of attending is perceived to be greater than the possible risk of infection with COVID-19 caused by leaving home. Added also emphasised advice to contact maternity services if concerns during pregnancy.
<b>7</b>	9.4.20	<b>3.1:</b> New section of reducing the risk to women of new infection caused by attending maternity settings. All other subsections in section 3 have been re-numbered.
<b>7</b>	9.4.20	<b>3.2:</b> New comment on visitor restrictions in maternity settings.
<b>7</b>	9.4.20	<b>3.2:</b> List of risk factors which contribute to mental ill health in pregnant women, and acknowledgement of the risk of increasing domestic violence with policy for social distancing, moved to section 3.6 on maternal mental wellbeing.
<b>7</b>	9.4.20	<b>3.3:</b> Advice about induction of labour changed to reference update to Saving Babies' Lives Care Bundle.
<b>7</b>	9.4.20	<b>4.2</b> Section 4.2 renamed 'Women with unconfirmed COVID-19 but symptoms suggestive of possible infection' to allow for inclusion of new recommendations on women who call the maternity unit with possible COVID-19 infection (not just attend in person).
<b>7</b>	9.4.20	<b>4.2:</b> Additional recommendations made to consider usual differential diagnoses in women who call the maternity unit to report a new fever/cough/respiratory symptoms.
<b>7</b>	9.4.20	<b>4.3.1:</b> New subsection added on the care of pregnant women who are self-isolating at home with suspected COVID-19.
<b>7</b>	9.4.20	<b>4.4:</b> Changed to subsection 4.3.3 (subsequent subsections re-numbered).
<b>7</b>	9.4.20	<b>4.6.1:</b> New recommendations re. prophylactic low molecular weight heparin to reduce risk of venous thromboembolism with COVID-19 infection in pregnancy, and to consider pulmonary embolism if women with COVID-19 suddenly deteriorate.
<b>7</b>	9.4.20	<b>4.7.2:</b> Statement on calling neonatal team early to inform them of imminent birth of a baby to a woman with COVID-19 moved to section 4.5, because it applies to all cases of COVID-19, not just in women with severe disease.

<b>8</b>	17.4.20	<b>1:</b> New paragraph on the quality of the available evidence and resultant classification of the advice.
<b>8</b>	17.4.20	<b>1.4:</b> New evidence included on the risk of COVID-19 in the woman, including a case series of pregnant women attending two maternity units in New York, who were screened for COVID-19 on arrival, the inclusion of the first report of maternal death directly attributed to COVID-19 in scientific literature and an update to the ICNARC data.
<b>8</b>	17.4.20	<b>4.2, 4.5.2 &amp; 4.6.2:</b> Restructured, including some new subtitles to organise and break up the text.
<b>8</b>	17.4.20	<b>4.3.1:</b> Renamed 'risk of venous thromboembolism'.
<b>8</b>	17.4.20	<b>4.6:</b> Section restructured for clarity.
<b>8</b>	17.4.20	<b>4.7 and 4.8:</b> Re-ordered the two sections within the text so that considerations for birth are written before considerations for neonatal and postnatal care.
<b>8</b>	17.4.20	<b>5.3:</b> Section re-structured. Also includes clarification that the recommendation for 10 days postnatal LMWH is regardless of mode of birth.
<b>8</b>	17.4.20	<b>Appendix 2:</b> Table of previous updates moved to appendix 3.
<b>8</b>	17.4.20	<b>Appendix 3:</b> New information on considerations when caring for women with suspected/confirmed COVID-19 during labour and birth.
<b>9</b>	13.5.20	<b>1:</b> Aims updated to include: The provision of safe, woman-centred care to women who are pregnant, give birth or are in the early postnatal period during the COVID-19 pandemic.
<b>9</b>	13.5.20	<b>1:</b> Findings of UKOSS data included in the summaries on viral transmission, effects on the woman and effects on the fetus/neonate. Where this supersedes existing references because of higher quality research or larger numbers, it has been used to replace it.
<b>9</b>	13.5.20	<b>1.3:</b> Updated information on possibility of vertical transmission to state that there are serious limitations to the available evidence.
<b>9</b>	13.5.20	<b>1.4:</b> Updated with emerging evidence on increased risk from COVID-19 to individuals with black, Asian and minority ethnic (BAME) background.
<b>9</b>	13.5.20	<b>2:</b> Information to share with pregnant women and their families has been removed from the guidance. All this information is also available in the RCOG information for pregnant women and their families in the COVID-19 hub. All subsequent sections have been renumbered.
<b>9</b>	13.5.20	<b>3.1 (Now 2.1):</b> Added paragraph about reducing transmission between staff.
<b>9</b>	13.5.20	<b>3.2 (Now 2.2):</b> Statement and recommendations added:  Emerging evidence suggests that individuals of black and minority ethnic (BAME) background may be at higher risk of developing severe complications of COVID-19. This may equally apply to pregnant women. We therefore advise:  <ul style="list-style-type: none"> <li>• Women of BAME background should be opportunistically advised that they may be at higher risk of complications of COVID-19, and advised to seek help early if they are concerned about their health.</li> <li>• Clinicians should be aware of this increased risk, and have a lower threshold to review, admit and consider multidisciplinary escalation in women of BAME background.</li> </ul>

9	13.5.20	<b>2.2:</b> Removed statement that further guidance on remote consultations will be published soon, and provided reference to RCM/RCOG guidance on antenatal and postnatal care.
9	13.5.20	<b>2.3:</b> Changed the statement that units should consider reducing provision of induction of labour for indications that are not 'strictly necessary', to units should consider reducing induction of labour where this is not 'medically indicated'.
9	13.5.20	<b>3.3 (Now 2.3):</b> Reference to NHS England 'Clinical guide for the temporary reorganisation of intrapartum maternity care during the coronavirus pandemic' added.
9	13.5.20	<b>3.3 (Now 2.3):</b> Statement added: 'Care should be taken to maintain safe services which continue to offer women support and choice as far as possible at this time. In particular, women should continue to be encouraged to contact their maternity unit with concerns about their or their baby's wellbeing. Justification should be provided for any service rationalisation required.'
9	13.5.20	<b>3.3 (Now 2.3):</b> Statement added: 'When reorganising services, maternity units should be particularly cognisant of emerging evidence that black, Asian and minority ethnic group (BAME) individuals are at particular risk of developing severe and life-threatening COVID-19. There is already extensive evidence on the inequality of experience and outcomes for BAME women during pregnancy and birth in the UK. Particular consideration should be given to the experience of women of BAME background and of lower socioeconomic status, when evaluating the potential or actual impact of any service change.'
9	13.5.20	<b>4.6 (Now 3.6):</b> Recommendation to be aware that myocardial injury is common among individuals with COVID-19, and reference added to <a href="#">NICE Guidance on diagnosis of myocardial injury in patients with suspected or confirmed COVID-19</a> .
9	13.5.20	<b>4.8.1 (Now 3.8.1):</b> Reference added to <a href="#">Resuscitation Council guidance</a> on neonatal life support during the COVID-19 pandemic.
9	13.5.20	<b>4.5.2 (Now 3.5.2):</b> Care in labour: Risk of venous thromboembolism. Clarification added that all women with suspected or confirmed COVID-19 should be discharged with 10 days' supply of prophylactic LMWH.
9	13.5.20	<b>4.4 (Now 3.4):</b> Women who develop new symptoms of COVID-19 during admission: Statement added that prophylaxis for venous thromboembolism should be considered and prescribed unless contraindicated.
9	13.5.20	<b>4.6 (Now 3.6):</b> Title change from 'Additional considerations in women with moderate/severe symptoms' to 'Women with suspected or confirmed COVID-19 and moderate/severe symptoms', to reflect that this includes information relevant to pregnant women admitted with COVID-19 outside of obstetric services.
9	13.5.20	<b>4.6 (Now 3.6):</b> Recommendation added: 'Prophylaxis for venous thromboembolism should be prescribed during admission unless contraindicated. At the time of discharge from hospital following a period of care for confirmed COVID-19 infection, all women should be prescribed at least 10 days of prophylactic LMWH.' This is consistent with recommendations already made elsewhere in previous versions of this document.
9	13.5.20	<b>4.6 (Now 3.6):</b> Changed statement 'Consider bacterial infection if the white blood cell count is raised (lymphocytes usually normal or low with COVID-19) and commence antibiotics' to 'Bacterial infection is an important differential diagnosis to COVID-19 infection. We advise blood cultures and a low threshold for antibiotics at presentation, with early review and rationalisation of antibiotics if COVID-19 is confirmed.'
9	13.5.20	<b>3.6:</b> Statement added: 'A woman with moderate or severe COVID symptoms who happens to be pregnant but with no immediate pregnancy issue should be cared for by the same multidisciplinary team as a non-pregnant woman with additional input from the maternity team. The labour ward should not be the default location for all pregnant women.'

## Summary of updates - Version 10

The following Version 10 summary of changes includes an additional column to reflect significant restructure changes between version 9 and 10 of this guidance.

Date	Summary of changes		
	Summary of changes	Section content from v9	Location in v10 update
4.6.20	<b>Introduction</b>		Now incorporates the following sections from v9: <ul style="list-style-type: none"> <li>• Purpose and scope</li> <li>• Identification and assessment of evidence</li> <li>• Epidemiology</li> <li>• Transmission</li> <li>• Effect of COVID-19 on pregnant women</li> <li>• Risk factors for hospital admission with COVID-19</li> <li>• Effect of COVID-19 on the fetus</li> </ul>
4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>2.2.</b> General advice regarding the continued provision of antenatal and postnatal services	<b>2.1</b> What are the considerations for organisation of antenatal care during the COVID-19 pandemic?
4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>2.3</b> General advice regarding possible service modifications during COVID-19	<b>2.2</b> What are the considerations for antenatal appointments?
4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>2.6</b> Smoking cessation and carbon monoxide monitoring in pregnancy	<b>2.3</b> What are the considerations for antenatal appointments?
4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>2.5</b> Maternal mental wellbeing	<b>2.2</b> What are the considerations for antenatal appointments?
4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>3.1</b> General advice for services providing care to pregnant women with suspected or confirmed COVID-19, where hospital attendance is necessary	<b>2.3</b> How should women with suspected or confirmed COVID-19 needing hospital attendance or advice be cared for?

4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>3.2</b> Women with unconfirmed COVID-19 but symptoms suggestive of possible infection	<b>2.3</b> How should women with suspected or confirmed COVID-19 needing hospital attendance or advice be cared for?
4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>3.3.3</b> Attendance for unscheduled/urgent antenatal care in women with suspected or confirmed COVID-19	<b>2.3</b> How should women with suspected or confirmed COVID-19 needing hospital attendance or advice be cared for?
4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>4.1</b> Antenatal care for pregnant women following self-isolation for symptoms suggestive of COVID-19	<b>2.4</b> What are the considerations for antenatal care for women who have recovered from COVID-19?
4.6.20	<b>Antenatal care during the COVID-19 pandemic</b>	<b>4.2</b> Antenatal care for pregnant women following hospitalisation for confirmed COVID-19 illness	<b>2.4</b> What are the considerations for antenatal care for women who have recovered from COVID-19?
4.6.20	<b>Venous thromboembolism prevention</b>	<b>3.3.1</b> Risk of venous-thromboembolism	<b>3.1</b> How should prevention of venous thromboembolism during the COVID-19 pandemic be addressed?
4.6.20	<b>Venous thromboembolism prevention</b>	<b>3.4</b> Women who develop new symptoms of COVID-19 during admission (antenatal, intrapartum or postnatal)  Sentence on thromboprophylaxis	<b>3.1</b> How should prevention of venous thromboembolism during the COVID-19 pandemic be addressed?
4.6.20	<b>Labour and birth</b>	<b>2.4</b> General advice regarding intrapartum services	<b>4.4</b> What about birth partners during the COVID-19 pandemic?
4.6.20	<b>Labour and birth</b>	Not in version 9	<b>New section in version 10: 4.1</b> What are the considerations for labour and birth in asymptomatic women who test or have tested positive for SARS-CoV-2?

4.6.20	<b>Labour and birth</b>	<b>3.5</b> Women attending for intrapartum care with suspected or confirmed COVID-19	<p><b>4.2</b> How should a woman with suspected/confirmed COVID-19 be looked after in labour if they are symptomatic?</p> <p><b>4.5</b> What informed discussions should take place with women regarding timing and mode of birth during the COVID-19 pandemic?</p> <p><b>4.6</b> What are the specific considerations for labour analgesia or anaesthesia?</p>
4.6.20	<b>Labour and birth</b>	<b>3.7</b> Specific peri-operative advice for healthcare professionals caring for pregnant women with suspected/confirmed COVID-19 who require surgical intervention	<p><b>4.8</b> How should obstetric theatres be managed during the COVID-19 pandemic?</p> <p><b>4.7</b> What personal protective equipment is recommended when caring for women during labour and birth?</p>
4.6.20	<b>Postnatal</b>	<b>3.8</b> Neonatal care	<p><b>6.1</b> How should neonatal care for the baby be provided during the COVID-19 pandemic?</p> <p><b>6.2</b> What should parents/carers be advised regarding infant feeding during the COVID-19 pandemic?</p>
4.6.20	<b>Postnatal</b>	<b>4.3</b> Postnatal care for pregnant women immediately following hospitalisation for confirmed COVID-19 illness	<b>6.3</b> What are the considerations for postnatal care for women and babies following admission with COVID-19?



## Appendix 2: Key considerations when caring for women with suspected/confirmed COVID-19 during labour and birth

<b>Consideration:</b>	
<b>Setting for birth</b>	If homebirth or birth in a midwifery-led unit is planned, a discussion should be initiated with the woman regarding the potentially increased risk of fetal compromise in active phase of labour if symptomatic with SARS-CoV-2. <sup>60</sup> Attending an obstetric unit, where the baby can be monitored using continuous electronic fetal monitoring (EFM), should be recommended for birth.
<b>Timing for birth</b>	<p>A positive COVID-19 result in an otherwise well woman, when there is also no evidence of fetal compromise, is not an indication to expedite birth.</p> <p>Induction of labour (IOL) is associated with longer periods of inpatient stay than for spontaneous onset of labour.</p> <p>Review the indication for IOL and consider whether the likely benefits outweigh possible risks. Where possible, review the provision and possibility of outpatient IOL.</p> <p>For women who are currently in a period of self-isolation because of suspected COVID-19 in themselves or a household contact, an individual assessment should be made to determine whether it is safe to delay scheduled appointments for pre-operative care and elective caesarean birth, or IOL if planned to occur during their period of self-isolation.</p> <p>The individualised assessment should consider the urgency of the birth, and the risk of infectious transmission to other women, healthcare workers and, postnatally, to her baby.</p>
<b>Mode for birth</b>	<p>There is currently no evidence to favour one mode of birth over another in women who are SARS-CoV-2 positive, so mode of birth should be discussed with the woman, taking into consideration her preferences and any obstetric indications for intervention.</p> <p>Mode of birth should not be influenced by the presence of COVID-19, unless the woman's respiratory condition demands urgent intervention for birth.</p> <p>The use of birthing pools in hospital should be avoided in suspected or confirmed cases, given that SARS-CoV-2 has been identified in faeces and that commonly available personal protective equipment (PPE) is not waterproof.</p> <p>An individualised informed discussion and decision should be made regarding shortening the length of the second stage of labour with instrumental birth in a symptomatic woman who is becoming exhausted or hypoxic.</p> <p>In case of deterioration in the woman's symptoms, an individual assessment should be made regarding the risks and benefits of continuing the labour versus proceeding to emergency caesarean birth, if this is likely to assist efforts to resuscitate the woman.</p> <p>Donning PPE is time-consuming. For emergency caesarean births, this may impact on the decision to delivery interval but it must be done. Women and their families should be told early about this possible delay.</p>

<p><b>Birth partners</b></p>	<p>Women should be supported and encouraged to have a birth partner present with them during their labour and birth. Having a trusted birth partner present throughout labour and birth is known to make a significant difference to the safety and wellbeing of women in childbirth.</p> <p>At a minimum, one asymptomatic birth partner should be permitted to stay with the woman through labour and birth, unless the birth occurs under general anaesthetic.</p> <p>When a woman contacts the maternity unit in early labour, she should be asked whether she or her birth partner have had any symptoms which could suggest COVID-19 in the preceding 7 days. If her partner has had onset of symptoms in the last 7 days, the woman should be advised that her partner should not attend the unit with her and she should consider bringing another birth partner who is symptom-free. Explain the need to protect maternity staff and other women and families from the risk of infection.</p> <p>On attendance to the maternity unit, all birth partners should also be asked whether they have had any symptoms which could suggest COVID-19 in the preceding 7 days. If the onset of these symptoms was within the last 7 days, or symptoms are still present (other than persistent cough), they should be asked to leave the maternity unit immediately and self-isolate at home.</p> <p>Birth partners should be asked to remain by the woman's bedside, to not walk around the ward/hospital and to wash their hands frequently.</p> <p>We recommend that birth partners be given clear advance guidance on what is expected of them should they need to accompany the woman to the operating theatre – e.g. for caesarean birth. This is particularly important given the challenges of staff communication when wearing full PPE.</p> <p>Restrictions on other visitors should follow hospital policy.</p>
<p><b>Respect and consent</b></p>	<p>Women must still be able to make decisions about the care they receive in line with the principles of informed consent</p>
<p><b>Fetal surveillance</b></p>	<p>Discuss with women the options for fetal surveillance in labour in accordance with existing NICE guidelines.</p> <p>Recommend continuous EFM for women who are symptomatic of COVID-19.</p> <p>Current infection with SARS-CoV-2 is not a contraindication for application of a fetal scalp electrode or for fetal blood sampling.</p>
<p><b>Pain relief</b></p>	<p>There is no evidence that epidural or spinal analgesia or anaesthesia is contraindicated in the presence of coronaviruses.</p> <p>Epidural analgesia should therefore be recommended in labour, to women with suspected or confirmed COVID-19 to minimise the need for general anaesthesia if urgent intervention for birth is needed.</p> <p>Entonox should be used with a single-patient microbiological filter. This is standard issue throughout maternity units in the UK.</p> <p>There is no evidence that the use of Entonox is an aerosol-generating procedure (AGP).</p>

<p><b>Intrapartum care</b></p>	<p>When a woman with confirmed or suspected COVID-19 is admitted to the maternity suite, the following members of the multidisciplinary team should be informed: consultant obstetrician, consultant anaesthetist, midwife-in-charge, consultant neonatologist, neonatal nurse in charge and infection control team.</p> <p>Maternal observations and assessment should be continued as per standard practice, with the addition of hourly oxygen saturations.</p> <p>Aim to keep oxygen saturation above 94%, titrating oxygen therapy accordingly.</p> <p>If the woman develops a fever, investigate and treat as per <a href="#">RCOG guidance on sepsis in pregnancy</a>, but also consider active COVID-19 as a cause of sepsis and investigate according to <a href="#">PHE guidance</a>.</p> <p>Apply caution with intravenous fluid management. Given the association of COVID-19 with acute respiratory distress syndrome, women with moderate-to-severe symptoms of COVID-19 should be monitored using hourly fluid input/output charts.</p> <p>Efforts should be targeted towards achieving neutral fluid balance in labour, in order to avoid the risk of fluid overload.</p>
<p><b>Immediate neonatal care</b></p>	<p>The neonatal team should be given sufficient notice of the birth, to allow them to attend and don PPE before entering the room/theatre.</p> <p>Given a lack of evidence to the contrary, delayed cord clamping is still recommended following birth, provided there are no other contraindications. The baby can be cleaned and dried as normal, while the cord is still intact.</p>
<p><b>Infection control</b></p>	<p>On arrival to hospital, women with suspected/confirmed COVID-19 should immediately be escorted to an isolation room or cohort bay/ward, suitable for the majority of care during their hospital visit or stay.</p> <p>Isolation rooms or ward bays should ideally have a defined area for staff to put on and remove PPE, and suitable bathroom facilities.</p> <p>Further advice on care in isolation rooms and COVID-19 cohort bays is <a href="#">available from PHE</a>.</p> <p>Only essential staff should enter the room and visitors should be kept to a minimum.</p> <p>All non-essential items from the clinic/scan room should be removed prior to the woman's arrival.</p> <p>General advice from PHE on type and specification of PPE is available <a href="#">available here</a>.</p>

<b>Pain relief</b>	<p>When a woman with confirmed or suspected COVID-19 is admitted to the maternity suite, the following members of the MDT should be informed: consultant obstetrician, consultant anaesthetist, midwife-in-charge, consultant neonatologist, neonatal nurse in charge and infection control team.</p> <p>Maternal observations and assessment should be continued as per standard practice, with the addition of hourly oxygen saturations.</p> <p>Aim to keep oxygen saturation more than 94%, titrating oxygen therapy accordingly.</p> <p>If the woman develops a fever, investigate and treat as per RCOG guidance on sepsis in pregnancy, but also consider active COVID-19 as a cause of sepsis and investigate according to PHE guidance.</p> <p>Apply caution with IV fluid management. Given the association of COVID-19 with acute respiratory distress syndrome, women with moderate to severe symptoms of COVID-19 should be monitored using hourly fluid input/output charts.</p> <p>Efforts should be targeted towards achieving neutral fluid balance in labour, in order to avoid the risk of fluid overload.</p>
<b>Infection control</b>	<p>Particular advice from Public Health England on type and specification of PPE for different maternity settings is available as part of the <a href="#">table here</a>.</p> <p>All clinical areas used must be cleaned after use, as per <a href="#">health protection guidance</a>.</p>



Royal College of  
Obstetricians &  
Gynaecologists

# References

## 8. References

1. Royal College of Obstetricians & Gynaecologists. Educational and support resources for coronavirus (COVID-19) 2020 [Available from: <https://www.rcog.org.uk/en/guidelines-research-services/coronavirus-covid-19-pregnancy-and-womens-health/educational-and-support-resources-for-coronavirus-covid-19/> accessed 01 June 2020.
2. Scottish Intercollegiate Guidelines Network. Implementation Support. Implementation Support [Available from: <https://www.sign.ac.uk/implementation-support.html> accessed 15 July 2019.
3. World Health Organisation. Coronavirus disease (COVID-2019) situation reports 2020 [Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/> accessed 01 June 2020.
4. Centers for Disease Control and Prevention. Human Coronavirus Types 2020 [Available from: <https://www.cdc.gov/coronavirus/types.html> accessed 01 June 2020.
5. Docherty AB, Harrison EM, Green CA, et al. Features of 16,749 hospitalised UK patients with COVID-19 using the ISARIC WHO Clinical Characterisation Protocol. medRxiv 2020:2020.04.23.20076042. doi: 10.1101/2020.04.23.20076042
6. Smith V, Seo D, Warty R, et al. Maternal and neonatal outcomes associated with COVID-19 infection: A systematic review. PLOS ONE 2020;15(6):e0234187. doi: 10.1371/journal.pone.0234187
7. ICNARC. Report on COVID-19 in critical care 2020 [updated 27 March. Available from: <https://www.icnarc.org/Our-Audit/Latest-News/2020/04/10/Report-On-5578-Patients-Critically-Ill-With-Covid-19>.
8. Lamouroux A, Attie-Bitach T, Martinovic J, et al. Evidence for and against vertical transmission for SARS-CoV-2 (COVID-19). American journal of obstetrics and gynecology 2020 doi: 10.1016/j.ajog.2020.04.039
9. Wang C, Zhou YH, Yang HX, et al. Intrauterine vertical transmission of SARS-CoV-2: what we know so far. Ultrasound Obstet Gynecol 2020;n/a(n/a) doi: 10.1002/uog.22045
10. Dong L, Tian J, He S, et al. Possible Vertical Transmission of SARS-CoV-2 From an Infected Mother to Her Newborn. JAMA 2020 doi: 10.1001/jama.2020.4621
11. Zeng H, Xu C, Fan J, et al. Antibodies in Infants Born to Mothers With COVID-19 Pneumonia. JAMA 2020 doi: 10.1001/jama.2020.4861
12. Schwartz D, Dhaliwal A. Infections in pregnancy with COVID-19 and other respiratory RNA virus diseases are rarely, if ever, transmitted to the fetus: Experiences with coronaviruses, HPIL, hMPV RSV, and influenza. Archives of Pathology & Laboratory Medicine doi: 10.5858/arpa.2020-0211-SA
13. Knight M, Bunch K, Vousden N, et al. Characteristics and outcomes of pregnant women admitted to hospital with confirmed SARS-CoV-2 infection in UK: national population based cohort study. BMJ 2020;369:m2107. doi: 10.1136/bmj.m2107
14. Martins-Filho PR, Santos VS, Santos HP, Jr. To breastfeed or not to breastfeed? Lack of evidence on the presence of SARS-CoV-2 in breastmilk of pregnant women with COVID-19. Rev Panam Salud

Publica 2020;44:e59. doi: 10.26633/RPSP:2020.59

15. Mor G, Cardenas I. The immune system in pregnancy: a unique complexity. *Am J Reprod Immunol* 2010;63(6):425-33. doi: 10.1111/j.1600-0897.2010.00836.x

16. Public Health England. COVID-19: investigation and initial clinical management of possible cases 2020 [Available from: <https://www.gov.uk/government/publications/wuhan-novel-coronavirus-initial-investigation-of-possible-cases/investigation-and-initial-clinical-management-of-possible-cases-of-wuhan-novel-coronavirus-wn-cov-infection> accessed 05 March 2020.

17. Guan W-j, Ni Z-y, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine* 2020 doi: 10.1056/NEJMoa2002032

18. Breslin N, Baptiste C, Gyamfi-Bannerman C, et al. COVID-19 infection among asymptomatic and symptomatic pregnant women: Two weeks of confirmed presentations to an affiliated pair of New York City hospitals. *Am J Obstet Gynecol MFM* 2020:100118. doi: 10.1016/j.ajogmf.2020.100118 [published Online First: 9 April 2020]

19. Sutton D, Fuchs K, D'Alton M, et al. Universal Screening for SARS-CoV-2 in Women Admitted for Delivery. *New England Journal of Medicine* 2020;382(22):2163-64. doi: 10.1056/NEJMc2009316

20. Knight M, Bunch K, Tuffnell D, et al. Saving Lives, Improving Mothers' Care 2019 [Available from: <https://www.npeu.ox.ac.uk/downloads/files/mbrance-uk/reports/MBRRACE-UK%20Maternal%20Report%202019%20-%20WEB%20VERSION.pdf>.

21. Liu Y, Chen H, Tang K, et al. Clinical manifestations and outcome of SARS-CoV-2 infection during pregnancy. *Journal of Infection* 2020; Online doi: <https://doi.org/10.1016/j.jinf.2020.02.028>

22. Vallejo V, Ilagan JG. A Postpartum Death Due to Coronavirus Disease 2019 (COVID-19) in the United States. *Obstet Gynecol* 2020 doi: 10.1097/AOG.0000000000003950

23. Hantoushzadeh S, Shamshirsaz AA, Aleyasin A, et al. Maternal Death Due to COVID-19 Disease. *American journal of obstetrics and gynecology* 2020 doi: 10.1016/j.ajog.2020.04.030

24. Karami P, Naghavi M, Feyzi A, et al. Mortality of a pregnant patient diagnosed with COVID-19: A case report with clinical, radiological, and histopathological findings. *Travel Med Infect Dis* 2020:101665. doi: 10.1016/j.tmaid.2020.101665

25. Henderson J, Gao H, Redshaw M. Experiencing maternity care: the care received and perceptions of women from different ethnic groups. *BMC Pregnancy Childbirth* 2013;13(1):196. doi: 10.1186/1471-2393-13-196

26. Raleigh VS, Hussey D, Seccombe I, et al. Ethnic and social inequalities in women's experience of maternity care in England: results of a national survey. *J R Soc Med* 2010;103(5):188-98. doi: 10.1258/jrsm.2010.090460

27. Office for National Statistics. Coronavirus (COVID-19) related deaths by ethnic group, England and Wales: 2 March 2020 to 10 April 2020 2020 [Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/coronavirusrelateddeathsbyethnicgroupenglandandwales/2march2020to10april2020> accessed 2020 11 May.

28. Khunti K, Singh AK, Pareek M, et al. Is ethnicity linked to incidence or outcomes of covid-19? *BMJ* 2020;369:m1548. doi: 10.1136/bmj.m1548
29. Marik PE, Kory P, Varon J. Does vitamin D status impact mortality from SARS-CoV-2 infection? *Med Drug Discov* 2020;100041. doi: 10.1016/j.medidd.2020.100041
30. Mitchell F. Vitamin D and COVID-1: do deficient risk a poorer outcome? *The Lancet Diabetes and Endocrinology* 2020
31. Pal BR, Marshall T, James C, et al. Distribution analysis of vitamin D highlights differences in population subgroups: preliminary observations from a pilot study in UK adults. *J Endocrinol* 2003;179(1):119-29. doi: 10.1677/joe.0.1790119
32. Grant WB, Lahore H, McDonnell SL, et al. Evidence that Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths. *Nutrients* 2020;12(4):988. doi: 10.3390/nu12040988 [published Online First: 2 April]
33. National Health Service. Vitamin D 2017 [Available from: <https://www.nhs.uk/conditions/vitamins-and-minerals/vitamin-d/> accessed 01 June 2020.
34. National Institute for Health and Care Excellence. Diabetes in pregnancy: management from preconception to the postnatal period 2015 [Available from: [www.nice.org.uk/guidance/ng3](http://www.nice.org.uk/guidance/ng3) accessed 23 March 2020.
35. Docherty AB, Harrison EM, Green CA, et al. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. *BMJ* 2020;369:m1985. doi: 10.1136/bmj.m1985
36. Zhang JP, Wang YH, Chen LN, et al. [Clinical analysis of pregnancy in second and third trimesters complicated severe acute respiratory syndrome]. *Zhonghua Fu Chan Ke Za Zhi* 2003;38(8):516-20.
37. Dowswell T, Carroli G, Duley L, et al. Alternative versus standard packages of antenatal care for low-risk pregnancy. *Cochrane Database Syst Rev* 2015(7):CD000934. doi: 10.1002/14651858.CD000934.pub3
38. Knight M, Bunch K, Tuffnell D, et al. Saving Lives, Improving Mothers' Care. Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2014–16. 2018 [updated November.
39. UK Government. Staying alert and safe (social distancing) 2020 [Available from: <https://www.gov.uk/government/publications/staying-alert-and-safe-social-distancing/staying-alert-and-safe-social-distancing> accessed 01 June 2020.
40. Peahl AF, Smith RD, Moniz MH. Prenatal Care Redesign: Creating Flexible Maternity Care Models Through Virtual Care. *American journal of obstetrics and gynecology* 2020 doi: 10.1016/j.ajog.2020.05.029
41. NHS England. Clinical guide for the management of remote consultations and remote working in secondary care during the coronavirus pandemic 2020 [Available from: <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/03/C0044-Specialty-Guide-Virtual-Working-and-Coronavirus-27-March-20.pdf> accessed 27 May 2020.



42. UK Government. Guidance on shielding and protecting people who are clinically extremely vulnerable from COVID-19 2020 [updated 18 May. Available from: <https://www.gov.uk/government/publications/guidance-on-shielding-and-protecting-extremely-vulnerable-persons-from-covid-19/guidance-on-shielding-and-protecting-extremely-vulnerable-persons-from-covid-19> accessed 27 May 2020.
43. Public Health England. COVID-19: Infection, prevention and control guidance 2020 [Available from: <https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control/wuhan-novel-coronavirus-wn-cov-infection-prevention-and-control-guidance> accessed 05 January 2020.
44. NHS England. Face masks and coverings to be worn by all NHS hospital staff and visitors 2020 [Available from: <https://www.gov.uk/government/news/face-masks-and-coverings-to-be-worn-by-all-nhs-hospital-staff-and-visitors> accessed 15 June 2020.
45. Knight M, Bunch K, Tuffnell D, et al. Saving Lives, Improving Mothers' Care 2019 [Available from: <https://www.npeu.ox.ac.uk/downloads/files/mbrance-uk/reports/MBRRACE-UK%20Maternal%20Report%202019%20-%20WEB%20VERSION.pdf> accessed 27 May 2020.
46. Higginbottom GMA, Evans C, Morgan M, et al. Experience of and access to maternity care in the UK by immigrant women: a narrative synthesis systematic review. *BMJ Open* 2019;9(12):e029478. doi: 10.1136/bmjopen-2019-029478
47. Vardavas CI, Nikitara K. COVID-19 and smoking: A systematic review of the evidence. *Tob Induc Dis* 2020;18(March):20. doi: 10.18332/tid/119324
48. NHS England. Saving babies' lives care bundle Version 2: COVID-19 information 2020 [Available from: <https://www.england.nhs.uk/publication/saving-babies-lives-care-bundle-version-2-covid-19-information/> accessed 27 May 2020.
49. Wu Y, Zhang C, Liu H, et al. Perinatal depressive and anxiety symptoms of pregnant women along with COVID-19 outbreak in China. *American journal of obstetrics and gynecology* 2020 doi: 10.1016/j.ajog.2020.05.009 [published Online First: 10 May]
50. Corbett GA, Milne SJ, Hehir MP, et al. Health anxiety and behavioural changes of pregnant women during the COVID-19 pandemic. *Eur J Obstet Gynecol Reprod Biol* 2020 doi: 10.1016/j.ejogrb.2020.04.022
51. Saccone G, Florio A, Aiello F, et al. Psychological Impact of COVID-19 in pregnant women. *American journal of obstetrics and gynecology* 2020 doi: 10.1016/j.ajog.2020.05.003
52. Royal College of Midwives. Domestic Abuse 2020 [Available from: [https://www.rcm.org.uk/media/4067/identifying-caring-for-and-supporting-women-at-risk-of\\_victims-of-domestic-abuse-during-covid-19-v1\\_\\_13052020final.pdf](https://www.rcm.org.uk/media/4067/identifying-caring-for-and-supporting-women-at-risk-of_victims-of-domestic-abuse-during-covid-19-v1__13052020final.pdf) accessed 27 May 2020.
53. Fraser E. Impact of COVID-19 Pandemic on Violence against Women and Girls 2020 [updated 16 March. Available from: <https://www.svri.org/sites/default/files/vawg-helpdesk-284-covid-19-and-vawg.pdf> accessed 27 May 2020.
54. UK Government. Coronavirus (COVID-19): support for victims of domestic abuse 2020 [Available from: <https://www.gov.uk/government/publications/coronavirus-covid-19-and-domestic-abuse/coronavirus-covid-19-support-for-victims-of-domestic-abuse> accessed 01 June 2020.

55. Royal College of Midwives. Personal Protective Equipment: Know your rights Guidance from the Royal College of Midwives 2020 [Available from: <https://www.rcm.org.uk/media/4060/ppe-know-your-rights-may-2020.pdf> accessed 27 May 2020.
56. Swartz D, Graham A. Potential Maternal and Infant Outcomes from Coronavirus 2019-nCoV (SARS-CoV-2) Infecting Pregnant Women: Lessons from SARS, MERS, and Other Human Coronavirus Infections. *Viruses* 2020;1-16.
57. Alserehi H, Wali G, Alshukairi A, et al. Impact of Middle East Respiratory Syndrome coronavirus (MERS-CoV) on pregnancy and perinatal outcome. *BMC Infect Dis* 2016;16:105. doi: 10.1186/s12879-016-1437-y
58. Bremme KA. Haemostatic changes in pregnancy. *Best Pract Res Clin Haematol* 2003;16(2):153-68. doi: 10.1016/s1521-6926(03)00021-5
59. Royal College of Obstetricians & Gynaecologists. Reducing the Risk of Venous Thromboembolism during Pregnancy and the Puerperium 2015 [updated April. Available from: <https://www.rcog.org.uk/globalassets/documents/guidelines/gtg-37a.pdf>.
60. Bikdeli B, Madhavan MV, Jimenez D, et al. COVID-19 and Thrombotic or Thromboembolic Disease: Implications for Prevention, Antithrombotic Therapy, and Follow-up. *J Am Coll Cardiol* 2020 doi: 10.1016/j.jacc.2020.04.031 [published Online First: 15 April]
61. Chen H, Guo J, Wang C, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet* 2020;395(10226):809-15. doi: 10.1016/S0140-6736(20)30360-3
62. Zhu H, Wang L, Fang C, et al. Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Transl Pediatr* 2020;9(1):51-60. doi: 10.21037/tp.2020.02.06
63. National Institute for Health and Care Excellence. Intrapartum Care for healthy women and babies 2017 [Available from: <https://www.nice.org.uk/guidance/cg190> accessed 27 May 2020.
64. NHS England. Clinical guide for the temporary reorganisation of intrapartum maternity care during the coronavirus pandemic 2020 [updated 9 April. Available from: <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/04/C0241-specialty-guide-intrapartum-maternity-care-9-april-2020.pdf> accessed 27 May 2020.
65. Amirian ES. Potential fecal transmission of SARS-CoV-2: Current evidence and implications for public health. *Int J Infect Dis* 2020;95:363-70. doi: 10.1016/j.ijid.2020.04.057
66. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA* 2020;323(18):1843-44. doi: 10.1001/jama.2020.3786
67. Bohren MA, Hofmeyr GJ, Sakala C, et al. Continuous support for women during childbirth. *Cochrane Database Syst Rev* 2017;7(7):CD003766. doi: 10.1002/14651858.CD003766.pub6
68. Bohren MA, Berger BO, Munthe-Kaas H, et al. Perceptions and experiences of labour companionship: a qualitative evidence synthesis. *Cochrane Database Syst Rev* 2019;3(3):CD012449. doi: 10.1002/14651858.CD012449.pub2

69. Shakibazadeh E, Namadian M, Bohren MA, et al. Respectful care during childbirth in health facilities globally: a qualitative evidence synthesis. *BJOG : an international journal of obstetrics and gynaecology* 2018;125(8):932-42. doi: 10.1111/1471-0528.15015 [published Online First: 2017/11/09]
70. NHS England. Visiting healthcare inpatient settings during the COVID-19 pandemic. 2020 [Available from: <https://www.england.nhs.uk/coronavirus/publication/visitor-guidance/>] accessed 15 June 2020.
71. Yang Z, Liu Y. Vertical Transmission of Severe Acute Respiratory Syndrome Coronavirus 2: A Systematic Review. *Am J Perinatol* 2020(EFirst) doi: 10.1055/s-0040-1712161 [published Online First: 13.05.2020]
72. Morau E, Bouvet L, Keita H, et al. Anaesthesia and intensive care in obstetrics during the COVID-19 pandemic. *Anaesth Crit Care Pain Med* 2020 doi: 10.1016/j.accpm.2020.05.006
73. Royal College of Anaesthetists. Obstetric Anaesthesia Guidance 2020 [Available from: <https://icmanaesthesiacovid-19.org/obstetric-anaesthesia-guidance> accessed 27 May 2020.
74. Royal College of Obstetricians & Gynaecologists. Bacterial sepsis in pregnancy 2020 [Available from: [https://www.rcog.org.uk/globalassets/documents/guidelines/gtg\\_64a.pdf](https://www.rcog.org.uk/globalassets/documents/guidelines/gtg_64a.pdf) accessed 27 May 2020.
75. Yang H, Hu B, Zhan S, et al. Effects of SARS-CoV-2 infection on pregnant women and their infants: A retrospective study in Wuhan, China. *Arch Pathol Lab Med* 2020 doi: 10.5858/arpa.2020-0232-SA
76. Shi L, Wang Y, Yang H, et al. Laboratory Abnormalities in Pregnant Women with Novel Coronavirus Disease 2019. *Am J Perinatol* 2020 doi: 10.1055/s-0040-1712181
77. World Health Organisation. Clinical management of severe acute respiratory infection when novel coronavirus (nCoV) infection is suspected 2020 [Available from: [https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected) accessed 05 March 2020.
78. The National Institute for Health and Care Excellence. COVID-19 rapid guideline: critical care in adults 2020 [updated March. Available from: <https://www.nice.org.uk/guidance/ng159>.
79. Scottish Intercollegiate Guidelines Network. COVID-19 position statement: Maternal critical care provision 2020 [Available from: [https://www.sign.ac.uk/assets/sg\\_maternal\\_critical\\_care\\_provision.pdf](https://www.sign.ac.uk/assets/sg_maternal_critical_care_provision.pdf) accessed 01 June 2020.
80. McLaren RA, Jr, London V, Atallah F, et al. Delivery For Respiratory Compromise Among Pregnant Women With COVID-19. *American journal of obstetrics and gynecology* 2020 doi: 10.1016/j.ajog.2020.05.035
81. Mathilde G, Rolnik DL, Hoffman MK, et al. Should we stop aspirin prophylaxis in pregnant women diagnosed with COVID-19? *Ultrasound Obstet Gynecol* 2020 doi: 10.1002/uog.22063
82. SARS-CoV-2 Surveillance Group. Characteristics of SARS-CoV-2 patients dying in Italy. Report based on available data on April 13th, 2020. 2020 [Available from: [https://www.epicentro.iss.it/en/coronavirus/bollettino/Report-COVID-2019\\_13\\_april\\_2020.pdf](https://www.epicentro.iss.it/en/coronavirus/bollettino/Report-COVID-2019_13_april_2020.pdf).
83. National Institute for Health and Care Excellence. COVID-19 rapid guideline: acute myocardial injury 2020 [updated 23 April 2020. Available from: <https://www.nice.org.uk/guidance/ng171> accessed 27 April 2020.

84. National Institute for Health and Care Excellence. Preterm labour and birth 2019 [updated August. Available from: <https://www.nice.org.uk/guidance/ng25> accessed 23 March 2020.
85. McIntosh JJ. Corticosteroid Guidance for Pregnancy during COVID-19 Pandemic. *Am J Perinatol* 2020(EFirst) doi: 10.1055/s-0040-1709684 [published Online First: 09.04.2020]
86. Thomson AJ, Royal College of O, Gynaecologists. Care of Women Presenting with Suspected Preterm Prelabour Rupture of Membranes from 24(+0) Weeks of Gestation: Green-top Guideline No. 73. *BJOG : an international journal of obstetrics and gynaecology* 2019;126(9):e152-e66. doi: 10.1111/1471-0528.15803
87. Royal College of Anaesthetists. Care of the critically ill woman in childbirth; enhanced maternal care 2018 [updated August 2018. Available from: <https://www.rcoa.ac.uk/sites/default/files/documents/2019-09/EMC-Guidelines2018.pdf> accessed 27 May 2020.
88. Royal College of Physicians. Acute care toolkit 15: Managing acute medical problems in pregnancy 2019 [updated October 2019. Available from: <https://www.rcplondon.ac.uk/guidelines-policy/acute-care-toolkit-15-managing-acute-medical-problems-pregnancy> accessed 27 May 2020.
89. University of Oxford. RECOVERY trial, interim results. 2020 [Available from: <https://www.recoverytrial.net/results>] accessed 16 June 2020.
90. UK Government. World first coronavirus treatment approved for NHS use by government. 2020 [Available from: <https://www.gov.uk/government/news/world-first-coronavirus-treatment-approved-for-nhs-use-by-government>] accessed 16 June 2020.
91. Chappell L, Williamson C, Knight M. Randomised Evaluation of COVID-19 Therapy (RECOVERY): Adaptation of trial protocol for pregnancy. 2020 [Available from: <https://www.recoverytrial.net/files/recovery-information-for-pregnant-patients-v5-0.pdf>] accessed 16 June 2020.
92. Royal College of Paediatrics and Child Health. COVID-19 - guidance for paediatric services 2020 [Available from: <https://www.rcpch.ac.uk/resources/covid-19-guidance-paediatric-services> accessed 01 June 2020.
93. Stuebe A. Should Infants Be Separated from Mothers with COVID-19? First, Do No Harm. *Breastfeed Med* 2020;15(5):351-52. doi: 10.1089/bfm.2020.29153.ams
94. Royal College of Midwives. Optimising mother-baby contact and infant feeding in a pandemic. Rapid analytic review 2020 [Available from: <https://www.rcm.org.uk/media/4096/optimising-infant-feeding-and-contact-rapid-review-19th-may-2020-submitted.pdf> accessed 27 May 2020.
95. Williams J, Namazova-Baranova L, Weber M, et al. The importance of continuing breastfeeding during COVID-19: in support to the WHO statement on breastfeeding during the pandemic. *J Pediatr* 2020 doi: 10.1016/j.jpeds.2020.05.009
96. UNICEF Baby Friendly Initiative. Statements on supporting infant feeding during the coronavirus (Covid-19) outbreak 2020 [Available from: <https://www.unicef.org.uk/babyfriendly/infant-feeding-during-the-covid-19-outbreak/> accessed 27 April 2020.
97. UK Government. Public advised to cover faces in enclosed spaces. 2020 [Available from: <https://www.gov.uk/government/news/public-advised-to-cover-faces-in-enclosed-spaces>] accessed 15 June 2020.

DISCLAIMER: The Royal College of Obstetricians and Gynaecologists (RCOG) has produced this guidance as an aid to good clinical practice and clinical decision-making. This guidance is based on the best evidence available at the time of writing, and the guidance will be kept under regular review as new evidence emerges. This guidance is not intended to replace clinical diagnostics, procedures or treatment plans made by a clinician or other healthcare professional and RCOG accepts no liability for the use of its guidance in a clinical setting. Please be aware that the evidence base for COVID-19 and its impact on pregnancy and related healthcare services is developing rapidly and the latest data or best practice may not yet be incorporated into the current version of this document. RCOG recommends that any departures from local clinical protocols or guidelines should be fully documented in the patient's case notes at the time the relevant decision is taken.

 @RCObsGyn    @rcobsgyn    @RCObsGyn



Royal College of  
Obstetricians &  
Gynaecologists

Royal College of Obstetricians and Gynaecologists, 10-18 Union Street, London, SE1 1SZ

T: +44 (0) 20 7772 6200

E: [covid-19@rcog.org.uk](mailto:covid-19@rcog.org.uk)

W: [rcog.org.uk](http://rcog.org.uk)

Registered Charity No. 213280