The routine immunisation schedule from Autumn 2018 **Diseases protected against** Age due Vaccine given and trade name **Usual site** Diphtheria, tetanus, pertussis (whooping cough), polio, Infanrix hexa Thigh DTaP/IPV/Hib/HepB Haemophilus influenzae type b (Hib) and hepatitis B Eight weeks old Pneumococcal Pneumococcal (13 serotypes) Prevenar 13 Thigh conjugate vaccine (PCV) Left thigh Meningococcal group B (MenB) MenB Bexsero Rotavirus gastroenteritis Rotavirus Rotarix By mouth Diphtheria, tetanus, pertussis, polio, DTaP/IPV/Hib/HepB Infanrix hexa Thigh Hib and hepatitis B Twelve weeks old Rotavirus Rotarix By mouth Rotavirus Diphtheria, tetanus, pertussis, polio, DTaP/IPV/Hib/HepB Infanrix hexa Thigh Hib and hepatitis B Sixteen weeks old Pneumococcal (13 serotypes) **PCV** Prevenar 13 Thigh Left thigh MenB MenB Bexsero Hib/MenC Upper arm/thigh Hib and MenC Menitorix **PCV** Pneumococcal Prevenar 13 Upper arm/thigh One year old (on or after the Measles, mumps and rubella MMR VaxPRO2 or **MMR** Upper arm/thigh child's first birthday) (German measles) **Priorix** MenB MenB booster Bexsero Left thigh Influenza (each year from Live attenuated Eligible paediatric age Fluenz Tetra² Both nostrils September) influenza vaccine LAIV³ groups1 Diphtheria, tetanus, pertussis and Infanrix IPV or DTaP/IPV Upper arm Three years four Repevax months old or soon MMR (check first MMR VaxPRO2 or after Measles, mumps and rubella Upper arm dose given) Priorix Cervical cancer caused by human Girls aged 12 to papillomavirus (HPV) types 16 and HPV (two doses Gardasil Upper arm 18 (and genital warts caused by 13 years 6-24 months apart) types 6 and 11) Td/IPV (check MMR Tetanus, diphtheria and polio Revaxis Upper arm status) Fourteen years old (school year 9) Meningococcal groups A, C, W and MenACWY Nimenrix or Menveo Upper arm Y disease Pneumococcal Pneumococcal Pneumococcal (23 serotypes) Polysaccharide Vaccine Polysaccharide Upper arm 65 years old (PPV) Vaccine Influenza (each year from Inactivated influenza 65 years of age Multiple Upper arm and older September) vaccine 70 years old Shingles Shingles Zostavax² Upper arm

2. Contains porcine gelatine.

All vaccines can be ordered from www.immform.dh.gov.uk free of charge except influenza for adults and pneumococcal polysaccharide vaccine.





^{1.} See Green book chapter 19 or visit www.gov.uk/government/publications/influenza-the-green-book-chapter-19 or www.nhs.uk/conditions/vaccinations/child-flu-vaccine/

^{3.} If LAIV (live attenuated influenza vaccine) is contraindicated and child is in a clinical risk group, use inactivated flu vaccine.

Selective imp	nunisation	program	mes

Target group	Age and schedule	Disease	Vaccines required
Babies born to hepatitis B infected mothers	At birth, four weeks and 12 months old ^{1,2}	Hepatitis B	Hepatitis B (Engerix B/HBvaxPRO)
Infants in areas of the country with TB incidence >= 40/100,000	At birth	Tuberculosis	BCG
Infants with a parent or grandparent born in a high incidence country ³	At birth	Tuberculosis	BCG
At risk children	From 6 months to 17 years of age	Influenza	Inactivated flu vaccine
Pregnant women	During flu season At any stage of pregnancy	Influenza	Inactivated flu vaccine
Pregnant women	From 16 weeks gestation	Pertussis	dTaP/IPV (Boostrix-IPV or Repevax)

- 1. Take blood for HBsAg at 12 months to exclude infection.
- In addition hexavalent vaccine (Infanrix hexa) is given at 8, 12 and 16 weeks.
 Where the annual incidence of TB is >= 40/100,000 see www.gov.uk/government/publications/tuberculosis-tb-by-country-rates-per-100000-people

Additional vaccines for individuals with underlying medical conditions

Medical condition	Diseases protected against	Vaccines required ¹	
Asplenia or splenic dysfunction (including due to sickle cell and coeliac disease)	Meningococcal groups A, B, C, W and Y Pneumococcal Haemophilus influenzae type b (Hib) Influenza	Hib/MenC MenACWY MenB PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine	
Cochlear implants	Pneumococcal	PCV13 (up to two years of age) PPV (from two years of age)	
Chronic respiratory and heart conditions (such as severe asthma, chronic pulmonary disease, and heart failure)	Pneumococcal Influenza	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine	
Chronic neurological conditions (such as Parkinson's or motor neurone disease, or learning disability)	Pneumococcal Influenza	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine	
Diabetes	Pneumococcal Influenza	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine	
Chronic kidney disease (CKD) (including haemodialysis)	Pneumococcal (stage 4 and 5 CKD) Influenza (stage 3, 4 and 5 CKD) Hepatitis B (stage 4 and 5 CKD)	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine Hepatitis B	
Chronic liver conditions	Pneumococcal Influenza Hepatitis A Hepatitis B	PCV13 (up to two years of age) PPV (from two years of age) Annual flu vaccine Hepatitis A Hepatitis B	
Haemophilia	Hepatitis A Hepatitis B	Hepatitis A Hepatitis B	
Immunosuppression due to disease or treatment ³	Pneumococcal Influenza	PCV13 (up to two years of age) ² PPV (from two years of age) Annual flu vaccine	
Complement disorders (including those receiving complement inhibitor therapy)	Meningococcal groups A, B, C, W and Y Pneumococcal <i>Haemophilus influenzae</i> type b (Hib) Influenza	Hib/MenC MenACWY MenB PCV13 (to any age) PPV (from two years of age) Annual flu vaccine	

- 1. Check relevant chapter of green book for specific schedule.
- 2. To any age in severe immunosuppression.3. Consider annual influenza vaccination for household members and those who care for people with these conditions.



